

STEDMAN'S Medical Dictionary

27th Edition

Illustrated in Color

STEDMAN'S Medical Dictionary

27th Edition

Illustrated in Color



LIPPINCOTT WILLIAMS & WILKINS

A Wolters Kluwer Company

Philadelphia • Baltimore • New York • London
Buenos Aires • Hong Kong • Sydney • Tokyo

Senior Managing Editor: Maureen Barlow Pugh
Managing Editor: Barbara Werner
New Terms Editor: Thomas W. Filardo, MD
Copy Editors: Peter W. Binns, Linda G. Francis, Raymond Lukens, Bonnie Montgomery
Chief On-Line Editor: Barbara L. Ferretti
On-Line Editors: Kathryn J. Cadle, Dana Workman
Proofreaders: Peter W. Binns; David A. Bloom, MD; Alfred J. Bollet, MD; Ted Burk; Regina Lavette Davis; John A. Day, Jr., MD, FCCP; Richard Diamanti; John H. Dirckx, MD; Thomas W. Filardo, MD; Linda G. Francis; John M. Last, MD, FRACP, FRCPC; Raymond Lukens; Kate Mason, CMT; Joan Sarchese
Database Programmers: Dave Marcus, Lexi-Comp Inc., Hudson, OH
Art Director: Jonathan Dimes
Illustrations: Neil O. Hardy
Additional artwork by: Mary Anna Barratt-Dimes, Kathryn Born, Rob Duckwall, Timothy Hengst, Mikki Senkarik, Michael Schenk, Larry Ward
Graphic preparation assistance: Susan Caldwell, Jennifer Clements, Thomas Dolan, Christina Nihira
Design: Dan Pfisterer

Copyright © 2000 Lippincott Williams & Wilkins
351 West Camden Street
Baltimore, Maryland 21201-2436 USA



Copyright © by William Wood and Company: 1911, 1st ed.; 1912, 2nd ed.; 1914, 3rd ed.; 1916, 4th ed.; 1918, 5th ed.; 1920, 6th ed.; 1922, 7th ed.; 1924, 8th ed.; 1926, 9th ed.; 1928, 10th ed.; 1930, 11th ed.

Copyright © by Williams & Wilkins: 1933, 12th ed.; 1935, 13th ed.; 1939, 14th ed.; 1942, 15th ed.; 1946, 16th ed.; 1949, 17th ed.; 1953, 18th ed.; 1957, 19th ed.; 1961, 20th ed.; 1966, 21st ed.; 1972, 22nd ed.; 1976, 23rd ed.; 1982, 24th ed.; 1990, 25th ed.; 1995, 26th ed.

All rights reserved. This book is protected by copyright. No part of this book may be reproduced in any form or by any means, including photocopying, or utilized by any information storage and retrieval system without written permission from the copyright owner.

Stedman's is a registered trademark of Lippincott Williams & Wilkins.

The publisher is not responsible (as a matter of product liability, negligence or otherwise) for any injury resulting from any material contained herein. This publication contains information relating to general principles of medical care which should not be construed as specific instructions for individual patients. Manufacturers' product information and package inserts should be reviewed for current information, including contraindications, dosages and precautions.

Database design by Lexi-Comp Inc., Hudson, OH
Printed in the United States of America by World Color, Inc.

Library of Congress Cataloging-in-Publication Data

Stedman, Thomas Lathrop, 1853-1938.
Stedman's medical dictionary.—27th ed.
p.; cm.
ISBN 0-683-40007-X (regular)—ISBN 0-683-40008-8 (deluxe)
1. Medicine—Dictionaries. I. Title: Medical dictionary. II. Title.
[DNLM: 1. Medicine—Dictionary—English. W 13 S812m 1999]
R121 .S8 1999
610'.3—dc21

99-056094

00 01 02 03 04 05
1 2 3 4 5

10

uremia (UO₂), a chronic metabolic disorder in which utilization of oxygen is decreased and the rate of carbohydrate, lipid, and protein metabolism is impaired and that of lipid and protein metabolism is decreased. It is caused by an absolute or relative deficiency of urea and is characterized, in more severe cases, by chronic azotemia, glycosuria, water and electrolyte loss, ketoacidosis, and long-term complications include neuropathy, renal osteodystrophy, generalized degenerative changes in large blood vessels, and increased susceptibility to infection. (Stained with honey)

DM is often associated with other risk factors, including obesity, hypertension, and disorders of lipid metabolism, obesity, hypercortisolemia, and insulin resistance, all of which are at least partially under genetic control. DM is often associated with other risk factors, including obesity, hypertension, and disorders of lipid metabolism, obesity, hypercortisolemia, and insulin resistance, all of which are at least partially under genetic control.

[illegible][illegible]

antenatal d., SYN prenatal d.

clinical d., a d. made from a study of the signs and symptoms of a disease.

differential d., the determination of which of two or more diseases with similar symptoms is the one from which the patient is suffering, by a systematic comparison and contrasting of the clinical findings. SYN differentiation (2).

d. by exclusion, a d. made by excluding those diseases to which only some of the patient's symptoms might belong, leaving one disease as the most likely d., although no definitive tests or findings establish that d.

laboratory d., a d. made by a chemical, microscopic, microbiologic, immunologic, or pathologic study of secretions, discharges, blood, or tissue.

neonatal d., systematic evaluation of the newborn for evidence of disease or malformations, and the conclusion reached.

pathologic d., a d., sometimes postmortem, made from an anatomic and/or histologic study of the lesions present.

physical d., (1) a d. made by means of physical examination of the patient. (2) the process of a physical examination.

prenatal d., d. utilizing procedures available for the recognition of diseases and malformations *in utero*, and the conclusion reached. SYN antenatal d.

di-ag-nos-tic (di-ag-nos'tik). 1. Relating to or aiding in diagnosis. 2. Establishing or confirming a diagnosis.

di-ag-nos-ti-cian (di-ag-nos-tish'än). One who is skilled in making diagnoses; formerly, a name for specialists in internal medicine.

Diagnostic and Statistical Manual of Mental Disorders (DSM). A system of classification, published by the American Psychiatric Association, that divides recognized mental disorders into clearly defined categories based on sets of objective criteria. Representing a majority view (rather than a consensus) of hundreds of contributors and consultants, DSM is widely recognized as a diagnostic standard and widely used for reporting, coding, and statistical purposes.


The first edition (1952), based on the sixth revision of the *International Classification of Diseases (ICD-6)*, was intended to promote uniformity in the naming and reporting of psychiatric disorders. It contained definitions of all named disorders, but no sets of diagnostic criteria. While its classification of mental disorders showed the influence of Freudian psychoanalysis, its nomenclature (e.g., depressive reaction, anxiety reaction, schizophrenic reaction) reflected the theories of Adolf Meyer (1866-1950). The second edition (*DSM-II*, 1968) preserved the psychoanalytic orientation but dropped the "reaction" terminology. The third edition (*DSM-III*, 1980) abandoned much of the rigidly psychodynamic thinking of the earlier editions and, for the first time, provided explicit diagnostic criteria and introduced a multiaxial system whereby different aspects of a patient's condition could be separately assessed. Briefly stated, the axes are I, clinical disorders; II, personality disorders and mental retardation; III, general medical disorders; IV, psychosocial and environmental stressors; and V, overall level of functioning. A revised version of the third edition (*DSM-III-R*, 1987) incorporated a number of improvements and clarifications. The fourth edition (*DSM-IV*) appeared in May, 1994. It follows its two predecessors closely in general outline, and like them is coordinated with and partly derived from *ICD-9*. For many observers, the most significant change in *DSM-IV* is the renaming of the category formerly called "Organic Mental Syndromes and Disorders" as "Delirium, Dementia, and Amnesic and Other Cognitive Disorders," a shift in terminology intended to avoid the implication that mental disorders in other categories are not organic.

di-a-gram. A simple, graphic depiction of an idea or object.

Dieuade d., SYN triaxial reference system.

flow d., a d. composed of blocks connected by arrows representing steps in a process such as decision analysis.

Venn d., pictorial representation of the extent to which two or more quantities or concepts are mutually inclusive and exclusive.



Venn diagram

di-a-ki-ne-sis (di'ä-ki-nē'sis). Final stage of prophase in meiosis I, in which the chiasmata present during the diplotene stage disappear, the chromosomes continue to shorten, and the nucleolus and nuclear membrane disappear. [G. *dia*, through, + *kinēsis*, movement]

dial (di'äl, dil). A clock face or instrument resembling a clock face. [L. *dies*, day]

astigmatic d., a diagram of radiating lines, used to test for astigmatism.

Di-a-lis-ter (di-äl-is'ter). An obsolete name for a genus of bacteria, the type species of which, *D. pneumosintes*, is now placed in the genus *Bacteroides*.

di-al-yl (di-al'il). A compound containing two allyl groups.

di-al-y-sance (di-al'i-sans). The number of milliliters of blood completely cleared of any substance by an artificial kidney or by peritoneal dialysis in a unit of time; conventional clearance formulas are expressed as mm/min. [fr. dialysis]

di-al-y-sate (di-al'i-sät). That part of a mixture that passes through a dialyzing membrane; the material that does not pass through is referred to as the retentate. SYN diffusate.

di-al-y-sis (di-al'i-sis). 1. A form of filtration to separate crystalline from colloid substances (or smaller molecules from larger ones) in a solution by interposing a semipermeable membrane between the solution and dialyzing fluid; the crystalline (smaller) substances pass through the membrane into the dialyzing fluid on the other side, the colloids do not. 2. The separation of substances across a semipermeable membrane on the basis of particle size and/or concentration gradients. 3. A method of artificial kidney function. [G. a separation, fr. *dialyo*, to separate]

continuous ambulatory peritoneal d. (CAPD), method of peritoneal d. performed in ambulatory patients with influx and efflux of dialysate during normal activities.

equilibrium d., in immunology, a method for determination of association constants for hapten-antibody reactions in a system in which the hapten (dialyzable) and antibody (nondialyzable) solutions are separated by semipermeable membranes. Since at equilibrium the quantity of free hapten will be the same in the two compartments, quantitative determinations can be made of hapten-bound antibody, free antibody, and free hapten.

extracorporeal d., hemodialysis performed through an apparatus outside the body.

peritoneal d., removal from the body of soluble substances and

water

which

perito

the bl

gradie

d. rel

senso

serrat

di-a-ly

from

di-a-ly

memt

di-a-m

magn

di-a-m

subst

ty, gi

pairec

conta

di-a-m

di-am

site p

body,

throug

measi

metro

anter

bipar

eminc

bucco

bucca

conju

conju

diago

exter

d. ob

obliq

sacro

syn d

obste

occip

occip

occip

poste

to the

right

subor

postel

fonta

total

tricle

tole.

total

cle in

trach

the ar

d. tri

trans

betw

zygo

arche

di-am

group

di-am

two e

di-a-n

amin

NH₂(

therapeutic i., the ratio of LD₅₀ to ED₅₀, used in quantitative comparison of drugs.

thoracic i., anteroposterior diameter of the thorax times 100 divided by the transverse diameter of the thorax. *syn* chest i.

tibiofemoral i., the ratio obtained by multiplying the length of the tibia by 100 and dividing by the length of the femur.

transversovertical i., *syn* vertical i.

tuberculoopsonic i., the opsonic i. calculated in relation to tuberculous infection, with an actively growing culture of *Mycobacterium tuberculosis* or the strain of tubercle bacillus from the patient being used in the test.

ultraviolet i., a daily i. issued by the U.S. National Weather Service for many cities, forecasting the amount of dangerous ultraviolet light that will arrive at the earth's surface about noon the following day.

uricolytic i., the percentage of uric acid oxidized to allantoin before being secreted.

vertical i., the relation of the height to the length of the skull: (height × 100)/length. *syn* height-length i., length-height i., transversovertical i.

vital i., the ratio of births to deaths within a population during a given time.

Volpe-Manhold i. (V-MI), an index for comparing the amount of dental calculus in individuals.

volume i., an indication of the relative size (e.g., volume) of erythrocytes, calculated as follows: hematocrit value, expressed as per cent of normal + red blood cell count, expressed as per cent of normal = volume i.

zygomatocauricular i., the ratio between the zygomatic and the auricular diameters of the skull or head.

in-di-can (in'di-kan). 1. Indoxyl β-D-glucoside from *Indigofera* species and *Polygonum tinctorium*; a source of indigo. *syn* plant i. 2. 3-Indoxylsulfuric acid, a substance found (as its salts) in sweat and in variable amounts in urine; indicative, when in quantity, of protein putrefaction in the intestine (indicanuria). *syn* metabolic i., uroanthin.

metabolic i., *syn* indican (2).

plant i., *syn* indican (1).

in-di-can-i-dro-sis (in'di-kan-i-drō'sis). Excretion of indican in the sweat. [indican + G. *hidrōs*, sweat]

in-di-cant (in'di-kant). 1. Pointing out; indicating. 2. An indication; especially a symptom indicating the proper line of treatment. [L. *in-dico*, pres. p. -ans (-ant), to point out]

in-di-can-u-ria (in'di-kan-ū'rē-ā). An increased urinary excretion of indican, a derivative of indol formed chiefly in the intestine when protein is putrefied; indol is also formed during the putrefaction of protein in other sites.

in-di-ca-tion (in-di-kā'shūn). The basis for initiation of a treatment for a disease or of a diagnostic test; may be furnished by a knowledge of the cause (causal i.), by the symptoms present (symptomatic i.), or by the nature of the disease (specific i.). [L. fr. *in-dico*, pp. -atus, to point out, fr. *dico*, to proclaim]

off label i., use of a medication for a purpose other than that approved by the FDA.

in-di-ca-tor (in'di-kā-ter, -tōr). 1. In chemical analysis, a substance that changes color within a certain definite range of pH or oxidation potential, or in any way renders visible the completion of a chemical reaction; e.g., litmus, phenolsulfonphthalein. 2. An isotope that is used as a tracer. 3. The labeled substance whose distribution between reactants of a system is used to determine the amount of analyte present. [L. one that points out]

alizarin i., a solution consisting of 1 g sodium alizarin sulfonate dissolved in 100 mL distilled water; used as an i. for free acidity in gastric contents.

clinical i., a measure, process, or outcome used to judge a particular clinical situation and indicate whether the care delivered was appropriate.

health i., variable, susceptible to direct measurement, that reflects the state of health of persons in a community.

oxidation-reduction i., a substance that undergoes a definite color change at a specific oxidation potential. *syn* redox i.

redox i., *syn* oxidation-reduction i.

in-di-ces (in'di-sēz). Alternative plural of index.

In-di-el-la (in-dē-el'ā). Old name for *Madurella*.

in-dig-e-nous (in-dij'ē-nūs). Native; natural to the country or region where found. [L. *indigenus*, born in fr. *indu*, within (old form of *in*), + G. -gen, producing]

in-di-ges-tion (in-di-jes'chūn). Nonspecific term for a variety of symptoms resulting from a failure of proper digestion and absorption of food in the alimentary tract.

acid i., i. resulting from hyperchlorhydria; often used by the laity as a synonym for pyrosis.

fat i., *syn* steatorrhea.

gastric i., *syn* dyspepsia.

nervous i., i. caused by emotional upsets or stress.

in-di-go (in'di-gō) [C.I. 73000]. A blue dyestuff obtained from *Indigofera tinctoria*, and other species of *Indigofera* (family Leguminosae); also made synthetically. *syn* indigo blue, indigotin. [L. *indicum*, fr. G. *indikon*, indigo, ntr. of *Indikos*, Indian]

in-di-go blue. *syn* indigo.

in-di-go car-mine [C.I. 73015]. A blue dye used for measurement of kidney function and as a special stain for Negri bodies. *syn* sodium indigotin disulfonate.

in-dig-o-tin (in-dig'ō-tin, in-di-gō'tin). *syn* indigo.

in-di-go-u-ria, in-di-gu-ria (in'di-gō-ū'rē-ā, in-di-goo'rē-ā). The excretion of indigo in the urine.

in-dis-po-si-tion (in-dis-pō-zish'ūn). Illness, usually slight; malaise. [L. *in* neg. + *dispositio*, an arrangement, fr. *dis-pono*, pp. -positus, to place apart]

in-di-um (In) (in'dē-ūm). A metallic element, atomic no. 49, atomic wt. 114.82. [*indigo*, because of its blue line in the spectrum]

in-di-um-111 (¹¹¹In). A cyclotron-produced radionuclide with a half-life of 2.8049 days and with gamma ray emissions of 171.2 and 245.3 kiloelectron volts. In a chloride form, it is used as a bone marrow and tumor-localizing tracer; in a chelate form, as a cerebrospinal fluid tracer. It is also used as a white blood cell labeling agent and as an antibody label.

i. chloride, i. trichloride, Cl₃In; used in electron microscopy to stain nucleic acids in thin tissue sections.

in-di-um-113m (^{113m}In). A radioactive isomer of ¹¹³In; it has a half-life of 1.658 hours; it has been used in cysternography and as a diagnostic aid in cardiac output.

in-di-vid-u-a-tion (in'di-vid-ū-ā'shūn). 1. Development of the individual from the specific. 2. In jungian psychology, the process by which one's personality is differentiated, developed, and expressed. 3. Regional activity in an embryo as a response to an organizer.

in-do-cy-a-nine green (in-dō-sī'ā-nēn). A tricarbocyanine dye that binds to serum albumin and is used in blood volume determinations and in liver function tests.

in-do-cy-bin (in-dō-sī'bin). *syn* psilocybin.

in-dol-ac-e-tu-ria (in'dōl-as-ē-too'rē-ā). Excretion of an appreciable amount of indoleacetic acid in the urine; a manifestation of Hartnup disease, also seen in patients with carcinoid tumors.

in-dol-a-mine (in-dōl'ā-mēn). General term for an indole or indole derivative containing a primary, secondary, or tertiary amine group (e.g., serotonin).

in-dole (in'dōl). 1. 2,3-Benzopyrrole; basis of many biologically active substances (e.g., serotonin, tryptophan); formed in degradation of tryptophan. *syn* ketole. 2. Any of many alkaloids containing the i. (1) structure.

in-do-lent (in'dō-lent). Inactive; sluggish; painless or nearly so, said of a morbid process. [L. *in*-neg. + *doleo*, pr. p. *dolens* (-ent), to feel pain]

in-dol-ic ac-ids (in-dōl'ik). Metabolites of L-tryptophan formed within the body or by intestinal microorganisms; the principal i. encountered in urine are indoleacetic acid, indoleacetylglutamine, 5-hydroxyindoleacetic acid, and indolelactic acid.

m. *flubis* *is* *pedis*, *official alternate term for lateral border of foot.
 m. *frontalis* *is* *palmaris*, *official alternate term for lateral border of palm.
 m. *frontalis* *ossis* *parietalis* [TA], *syn* frontal border of parietal bone.
 m. *frontalis* *ossis* *sphenoidalis* [TA], *syn* frontal margin of sphenoid.
 m. *lacinia* [TA], *syn* labial margin.
 m. *lateralis* [TA], *syn* inferolateral border.
 m. *lateralis* *cer* *ebri*, *syn* inferolateral margin of cerebellar hemisphere.
 m. *lateralis* *corporis* *pancreatis* [TA], *syn* lateral border of body of pancreas.
 m. *lateralis* *corporis* *spenis*, *syn* inferior border of body of penis.
 m. *lateralis* *hup* *is* *etis* [TA], *syn* inferior border of liver.
 m. *lateralis* *pancreatis* *is*, *syn* inferior border of body of pancreas.
 m. *lateralis* *pauis* *is* [TA], *syn* anterior border of lung.
 m. *lateralis* *spenis* [TA], *syn* inferior border of spleen.
 m. *lateralis* *vis* [TA], *syn* inferolateral margin of cerebellar hemisphere.
 m. *intermedia* *is* *hemispheri* *cerebri* [TA], *syn* inferomedial border of cerebellar hemisphere.
 m. *intermedia* *is*, *syn* inferomedial margin.
 m. *intermedia* *is*, *syn* inferolateral margin.
 m. *interna* *is* *vis* *fibulae* [TA], *syn* interosseous border of fibula.
 m. *interna* *is* *vis* *radius* [TA], *syn* interosseous border of radius.
 m. *interna* *is* *vis* *tibiae* [TA], *syn* interosseous border of tibia.
 m. *interna* *is* *vis* *of* *the* *TA*, *syn* interosseous border of tibia.
 m. *lacrime* *is* *maxillae* [TA], *syn* inferomedial margin of maxilla.
 m. *lumboides* *ossis* *occipitalis* [TA], *syn* lambdoid border of occipital bone.
 m. *lumboides* *ossis* *occipitalis* *is*, *syn* lambdoid border of occipital bone.
 m. *lateralis* [TA], *syn* lateral border.
 m. *lateralis* *is* *antebis* *chili*, *official alternate term for radial border of forearm.
 m. *lateralis* *is* *humeri* *is*, *syn* lateral border of humerus.
 m. *lateralis* *is* *pedis* [TA], *syn* lateral border of foot.
 m. *lateralis* *is* *re* *vis* [TA], *syn* lateral border of kidney.
 m. *lateralis* *is* *scapulae* *is*, *syn* lateral border of scapula.
 m. *lateralis* *is* *vis* *gulis* [TA], *syn* lateral border of nail.
 m. *liber* [TA], *syn* free border.
 m. *liber* *ova* *ri* [TA], *syn* free border of ovary.
 m. *liber* *vis* *gulis* [TA], *syn* free border of nail.
 m. *liber* *vis* *margin* *is*, *syn* margin of tongue.
 m. *nasoides* *ossis* *occipitalis* [TA], *syn* mastoid border of occipital bone.
 m. *nasoides* *ossis* *occipitalis* *is*, *syn* mastoid border of occipital bone.
 m. *media* [TA], *syn* medial border.
 m. *media* *is* *antebis* *chili*, *official alternate term for ulnar border of forearm.
 m. *media* *is* *cer* *ebri*, *syn* inferomedial margin of cerebellar hemisphere.
 m. *media* *is* *glan* *is* *vis* *supraure* *is* [TA], *syn* medial border of suprarenal gland.
 m. *media* *is* *humeri* *is*, *syn* medial border of humerus.
 m. *media* *is* *is* *pedis* [TA], *syn* medial border of foot.
 m. *media* *is* *re* *vis* [TA], *syn* medial border of kidney.
 m. *media* *is* *scapulae* *is*, *syn* medial border of scapula.
 m. *media* *is* *vis* *tibiae* [TA], *syn* medial border of tibia.
 m. *meso* *is* *fibula* *is*, *syn* mesoventral border of ovary.
 m. *nasoides* *is* *ossis* *frontalis* [TA], *syn* nasal margin of frontal bone.
 m. *occipitalis* [TA], *syn* occipital border.
 m. *occipitalis* *is* *os* *parietalis* [TA], *syn* occipital border of parietal bone.

[illegible][illegible]

tumor markers used in primary diagnoses

[illegible]

Markov, Andrei, Russian mathematician, 1865-1922. see *Mar-*
kov process.

Marmé re a gent. See under *reagent*.

mar-mar-ri-ated (mar'-mō-rī-ated). Denoting a condition in which the appearance of the skin is streaked like marble, see also *cuius mar-mar-riatus* [L. *mar-mar-riatus*, mottled].

mar-trot (mar'-trot). A woodcock or groundhog; aibernating rodent that may serve as reservoir host of plague bacillus in North America. [Fr. *martrot*].

Marotzau, Pierre, French medical geneticist. *1926. see *Mar-*
otzau syndrome.

Marquis re a gent. See under *reagent*.

mar-mar-qu-ri-ated (mar'-mā-rī-ated). 1. A highly cellular hematopoietic con-marrow (mar'-ō) [TA]. 2. A highly cellular hematopoietic con-marrow containing the medullary sinus and spongy epithyses of bone; it is located in the medullary cavity of the shaft of the long bones of the limbs. 2. Any soft gelatinous or fatty material resembling the m. of bone. see also *medulla*. [A.S. *marz*]

mar-row (mar'-ō) [TA]. The soft, pulpy tissue filling the medullary cavities of bones, having a spongy or reticular form and cells; it differs in consistency by age and location. see also *gelatinous bone m.*, *red bone m.*, *yellow bone m.* *syn medulla osseum* [TA].

gelatinous bone m. [TA]. Degenerated marrow of cranial bones in old age.

red bone m. [TA]. bone marrow in which the stroma primarily contain the developmental stages of erythrocytes, leukocytes, and megakaryocytes; it is present throughout the skeleton during fetal development. After the first postnatal year, it is gradually replaced in the long bones by yellow marrow. *syn medulla osseum rubra* [TA].

spinal m., *syn spinal cord*.

yellow marrow m. [TA]. bone m. in which the stroma of the reticu-lar network are largely filled primarily with fat; it replaces red marrow in the long bones after the fifth year of life. *syn medulla osseum flava* [TA].

Marshall, Dian K., U.S. ophthalmologist. *1905. see *M. syndrome*.

Marshall, Eli K., U.S. pharmacologist. 1898-1966. see *M. meth-*
os.

Marshall, John, English anatomist, 1818-1891. see *M. vestig-*
ial, oblique vein.

Marshall, Victor P., U.S. urologist. *1913. see *M. leri*; *M-*
Marchei test; *M.-Marchei-Krantz operation*.

Mar-shal-la-gi' mar-shāl-lī (mar-shā-lā'-mā-shāl-lī). One of the medium stomach worms of the nematode family Trichostrongyidae, found in the abomasum of sheep, goats, camels, and various wild ruminants.

marsh wall root (marsh'-wāl-ōt). *syn althaea*.

mar-sū-pū-al (mar'-sū-pē-ū-al). 1. A member of the order, *Mar-*
supula, which includes such mammals as kangaroos, wombats, bandicoots, and opossums; the female of which has an abdominal pouch for carrying the young. 2. Of or pertaining to marsupials.

mar-sū-pū-al-ia (mar'-sū-pē-ū-ā-lī-ā). Ectoparasitization of the *syn* or other insect infestation by feeding the superior edges of the skin, thereby causing a pouch. [L. *mar-sū-pū-al*, pouch]

mar-sū-pū-al-um (mar'-sū-pē-ū-ū-m). 1. *syn* scrotum. 2. A pouch or fold. [L. *mar-sū-pū-al*, pouch]

Mar-tel-lini, J., 19th century Italian anatomist. see *M. area*, *fund.*

Martin, August E., German gynecologist, 1847-1933. see *M-*
ube; *M.-Gruber anastomosis*.

Martini, Henry A., U.S. surgeon, 1824-1884. see *M. bandage*, *disease*.

Martin, J.E. see Thayer-M. medium.

Martino-tti, Giovanni (Italian physician, 1857-1928. see *M. cell*).

mar-tin-ut-us vel-mus (mar'-tī-nū-ū-tū-s) [C.L. 10313]. An acid dye used as a stain in plant and animal histology, and as a light filter for photo-microscopy. [Lat. *A. Martinus*, Chem. artist, *1920].

difficult for patients to make informed choices about their care. Patients should be told who is providing care, what benefits and burdens can be attributed to trainees, and how trainees are supervised. Most patients, when informed, allow trainees to play an active role in their care.

IMPAIRED PHYSICIANS Physicians may hesitate to intervene when colleagues impaired by alcohol abuse, drug abuse, or psychiatric or medical illness place patients at risk. However, society relies on physicians to regulate themselves. If colleagues of an impaired physician do not take steps to protect patients, no one else may be in a position to do so.

CONFLICTS FOR TRAINEES Medical students and residents may fear that they will receive poor grades or evaluations if they act on the patient's behalf by disclosing mistakes, avoiding misrepresentation of their role, and reporting impaired colleagues. Discussing such dilemmas with more senior physicians can help trainees check their interpretation of the situation and obtain advice and assistance.

ADDITIONAL ETHICAL ISSUES

MAINTAINING CONFIDENTIALITY Maintaining the confidentiality of medical information respects patients' autonomy and privacy, encourages them to seek treatment and to discuss their problems candidly, and prevents discrimination. Physicians need to guard against inadvertent breaches of confidentiality, as when talking about patients in elevators. Maintaining confidentiality is not an absolute rule. The law may require physicians to override confidentiality in order to protect third parties, for example, reporting to government officials persons with specified infectious conditions, such as tuberculosis and syphilis; persons with gunshot wounds; and victims of elder abuse and domestic violence. Computerized medical records raise additional concerns because breaches of confidentiality may affect many patients.

ALLOCATING RESOURCES JUSTLY Allocation of limited health care resources is problematic. Ideally, allocation decisions should be made as public policy, with physician input. At the bedside, physicians generally should act as patient advocates within constraints set by society, reasonable insurance coverage, and sound practice. *Ad hoc* rationing by the individual physician at the bedside may be inconsistent, discriminatory, and ineffective. In some cases, however, two patients may compete for the same limited resources, such as physician time or a bed in intensive care. When this occurs, physicians should ration their time and resources according to patients' medical needs and the probability of benefit.

ASSISTANCE WITH ETHICAL ISSUES Discussing perplexing ethical issues with other members of the health care team, colleagues, or the hospital ethics committee often clarifies issues and suggests ways to improve communication and to deal with strong emotions. When struggling with difficult ethical issues, physicians may need to reevaluate their basic convictions, tolerate uncertainty, and maintain their integrity while respecting the opinions of others.

BIBLIOGRAPHY

- ALPERS A, LO B: When is CPR futile? *JAMA* 273:156, 1995
- AMERICAN COLLEGE OF PHYSICIANS: *American College of Physicians Ethics Manual*. Ann Intern Med. 128:576, 1998
- BEAUCHAMP TL, CHILDRESS JP: *Principles of Biomedical Ethics*, 5th ed. New York, Oxford University Press, 2000
- EMANUEL EJ et al: The practice of euthanasia and physician-assisted suicide in the United States: Adherence to proposed safeguards and effects on physicians. *JAMA* 280:507, 1998
- GRISIO T, APPELBAUM P: *Assessing Competence to Consent to Treatment: A Guide for Physicians and Other Health Professionals*. New York, Oxford University Press, 1998
- KASSIRER JP: Managed care and the morality of the marketplace. *N Engl J Med* 333:50, 1995

LO B: *Resolving Ethical Dilemmas: A Guide for Clinicians*, 2d ed. Philadelphia, Lippincott Williams & Wilkins, 2000

MEISEL A: *The Right to Die*, 2d ed. New York, Wiley, 1995

QUILL TE et al: Palliative options of last resort: A comparison of voluntarily stopping eating and drinking; terminal sedation; physician-assisted suicide; and voluntary active euthanasia. *JAMA* 278:2099, 1997

TULSKY JA et al: Opening the black box: How do physicians communicate about advance directives. *Ann Intern Med* 129:441, 1998

3

Daniel B. Mark

DECISION-MAKING IN CLINICAL MEDICINE

To the medical student who requires 2 h to collect a patient's history and perform a physical examination, and several additional hours to organize them into a coherent presentation, the experienced clinician's ability to reach a diagnosis and decide on a management plan in a fraction of the time seems extraordinary. While medical knowledge and experience play a significant role in the senior clinician's ability to arrive at a differential diagnosis and plan quickly, much of the process involves skill in clinical decision-making. The first goal of this chapter is to provide an introduction to the study of clinical reasoning.

Equally bewildering to the student are the proper use of diagnostic tests and the integration of the results into the clinical assessment. The novice medical practitioner typically uses a "shotgun" approach to testing, hoping to hit a target without knowing exactly what that target is. The expert, on the other hand, usually has a specific target in mind and efficiently adjusts the testing strategy to it. The second goal of this chapter is to review briefly some of the crucial basic statistical concepts that govern the proper interpretation and use of diagnostic tests; quantitative tools available to assist in clinical decision-making will also be discussed.

CLINICAL DECISION-MAKING

CLINICAL REASONING The most important clinical actions are not procedures or prescriptions but the judgments from which all other aspects of clinical medicine flow. In the modern era of large randomized trials, it is easy to overlook the importance of this elusive mental activity and focus instead on the algorithmic practice guidelines constructed to improve care. One reason for this apparent neglect is that much more research has been done on how doctors *should* make decisions (e.g., using a Bayesian model discussed below) than on how they actually *do*. Thus, much of what we know about clinical reasoning comes from empirical studies of nonmedical problem-solving behavior.

Despite the great technological advances of the twentieth century, uncertainty still plays a pivotal role in all aspects of medical decision-making. We may know that a patient does not have long to live, but we cannot be certain how long. We may prescribe a potent new receptor blocker to reverse the course of a patient's illness, but we cannot be certain that the therapy will do so without side effects. Uncertainty in medical outcomes creates the need for probabilities and other mathematical/statistical tools to help guide decision-making. (These tools are reviewed later in the chapter.)

Uncertainty is compounded by the information overload that characterizes modern medicine. Today's experienced clinician needs close to 2 million pieces of information to practice medicine. Doctors subscribe to an average of 7 journals, representing over 2500 new articles each year. Computers offer the obvious solution both for management of information and for better quantitation and management of the daily uncertainties of medical care. While the technology to computerize medical practice is available, many practical problems remain to be solved before patient information can be standardized and integrated with medical evidence on a single electronic platform.

The following three examples introduce the subject of clinical reasoning:

A 46-year-old man presents to his internist with a chief complaint of hemoptysis. The physician knows that the differential diagnosis of hemoptysis includes over 100 different conditions, including cancer and tuberculosis (Chap. 33). The examination begins with some general background questions, and the patient is asked to describe his symptoms and their chronology. By the time the examination is completed, and even before any tests are run, the physician has formulated a working diagnostic hypothesis and planned a series of tests to test it. In an otherwise healthy and nonsmoking patient recovering from a viral bronchitis, the doctor's hypothesis would be that the acute bronchitis is responsible for the small amount of blood-streaked sputum the patient observed. In this case, a chest x-ray and purified protein derivative (PPD) skin test may be sufficient.

A second 46-year-old patient with the same chief complaint who had a 100-pack-year smoking history, a productive morning cough, and episodes of blood-streaked sputum may generate the principal diagnostic hypothesis of carcinoma of the lung. Consequently, along with the chest x-ray and PPD skin test, the physician refers this patient for bronchoscopy.

A third 46-year-old patient with hemoptysis who is from a developing country is evaluated with an electrocardiogram as well, because the physician thinks she has a soft diastolic rumble at the apex and cardiac auscultation, suggesting rheumatic mitral stenosis. These three vignettes illustrate two aspects of expert clinical reasoning: (1) the use of cognitive shortcuts, or *heuristics*, as a way to organize the complex, unstructured material that is collected in the clinical evaluation; and (2) the use of diagnostic hypotheses to consolidate the information and indicate appropriate management steps.

THE USE OF COGNITIVE SHORTCUTS. Heuristics reduce the complexity of a problem to a manageable level. Psychologists have found that people rely on three basic types of heuristics. For example, when assessing a patient, clinicians often weigh the probability that this patient's clinical features match those of the class of patients with the leading diagnostic hypothesis being considered. In other words, the clinician is searching for the diagnosis for which the patient appears to be a representative example; this cognitive shortcut is called the *representativeness heuristic*. It may take only a few characteristics from the history for an expert clinician using the representativeness heuristic to arrive at a sound diagnostic hypothesis. For example, an elderly patient with new-onset fever, cough, productive of copious sputum, unilateral pleuritic chest pain, and dyspnea is readily identified as fitting the pattern for acute pneumonia, probably of bacterial origin. Evidence of focal pulmonary consolidation on the physical examination will increase the clinician's confidence in the diagnosis because it fits the expected pattern of acute bacterial pneumonia. Knowing this allows the experienced clinician to conduct an efficient, directed, and therapeutically productive patient evaluation although there may be little else in the history or physical examination of direct relevance. The inexperienced medical student or resident, who has not yet learned the patterns most prevalent in clinical medicine, must work much harder to achieve the same result and is often at risk of missing the important clinical problem in a sea of comparatively collected but unhelpful data.

However, physicians using the representativeness heuristic can reach erroneous conclusions if they fail to consider the underlying prevalence of two competing diagnoses. Consider a patient with pleuritic chest pain, dyspnea, and a low-grade fever. A clinician might consider acute pneumonia and acute pulmonary embolism to be the two leading diagnostic alternatives. Clinicians using the representativeness heuristic might judge both diagnostic candidates to be equally likely, although to do so would be wrong if pneumonia was much more prevalent in the underlying population. Mistakes may also result from a failure to consider that a pattern based on a small number of

QUESTIONS: The Right to Die, 2d ed. New York, Wiley, 1995.
 QUESTIONS: The Right to Die, 2d ed. New York, Wiley, 1995.
 QUESTIONS: The Right to Die, 2d ed. New York, Wiley, 1995.
 QUESTIONS: The Right to Die, 2d ed. New York, Wiley, 1995.

DECISION-MAKING IN CLINICAL MEDICINE

Daniel B. Mark

To the medical student who requires 24 to collect a patient's history and perform a physical examination, and several additional hours to organize them into a coherent presentation, the experienced clinician's ability to reach a diagnosis and decide on a management plan in a fraction of the time seems extraordinary. While medical knowledge and experience play a significant role in the senior clinician's ability to arrive at a differential diagnosis and plan quickly, much of the process involves skill in clinical decision-making. The first goal of this chapter is to provide an introduction to the study of clinical reasoning. Equally bewildering to the student are the proper use of diagnostic tests and the integration of the results into the clinical assessment. The novice medical practitioner typically uses a "shotgun" approach to testing, hoping to hit a target without knowing exactly what that target is. The expert, on the other hand, usually has a specific target in mind and efficiently adjusts the testing strategy to it. The second goal of this chapter is to review briefly some of the crucial basic diagnostic concepts that govern the proper interpretation and use of diagnostic tests; quantitative tools available to assist in clinical decision-making will also be discussed.

CLINICAL REASONING

The most important clinical actions are not procedures or prescriptions but the judgments from which all other aspects of clinical medicine flow. In the modern era of large randomized trials, it is easy to overlook the importance of this elusive mental activity and focus instead on the algorithmic practice guidelines constructed to improve care. One reason for this apparent neglect is that much more research has been done on how doctors *should* make decisions (e.g., using a Bayesian model discussed below) than on how they actually *do*. Thus, much of what we know about clinical reasoning comes from empirical studies of nonmedical, problem-solving behavior.

Despite the great technological advances of the twentieth century, uncertainty still plays a pivotal role in all aspects of medical decision-making. We may know that a patient does not have long to live, but we cannot be certain how long. We may prescribe a potent new receptor blocker to reverse the course of a patient's illness, but we cannot be certain that the therapy will do so without side effects. Uncertainty in medical outcomes creates the need for probabilities and other mathematical/statistical tools to help guide decision-making. (These tools are reviewed later in the chapter.)

Uncertainty is compounded by the information overload that characterizes modern medicine. Today's experienced clinician needs close to 2 million pieces of information to practice medicine. Doctors subscribe to an average of 7 journals, representing over 2500 new articles each year. Computers offer the obvious solution both for management of information and for better quantitation and management of the daily uncertainties of medical care. While the technology to computerize medical practice is available, many practical problems remain to be solved before patient information can be standardized and integrated with medical evidence on a single electronic platform.

difficult for patients to make informed choices about their care. Patients should be told who is providing care, what benefits and burdens can be attributed to trainees, and how trainees are supervised. Most patients, when informed, allow trainees to play an active role in their care.

IMPAIRED PHYSICIANS. Physicians may hesitate to intervene when colleagues impaired by alcohol abuse, drug abuse, or psychiatric or medical illness place patients at risk. However, society relies on physicians to regulate themselves. If colleagues of an impaired physician do not take steps to protect patients, no one else may be in a position to do so.

CONFLICTS FOR TRAINERS. Medical students and residents may fear that they will receive poor grades or evaluations if they act on the patient's behalf by disclosing mistakes, avoiding misrepresentation of their role, and reporting impaired colleagues. Discussing such dilemmas with more senior physicians can help trainees check their interpretation of the situation and obtain advice and assistance.

ADDITIONAL ETHICAL ISSUES

MAINTAINING CONFIDENTIALITY. Maintaining the confidentiality of medical information respects patients' autonomy and privacy, encourages them to seek treatment and to discuss their problems candidly, and prevents discrimination. Physicians need to guard against inadvertent breaches of confidentiality, as when talking about patients in elevators. Maintaining confidentiality is not an absolute rule. The law may require physicians to override confidentiality in order to protect third parties, for example, reporting to government officials persons with specified infectious conditions, such as tuberculosis and syphilis; persons with gunshot wounds; and victims of elder abuse and domestic violence. Computerized medical records raise additional concerns because breaches of confidentiality may affect many patients.

ALLOCATING RESOURCES JUSTLY. Allocation of limited health care resources is problematic. Ideally, allocation decisions should be made as public policy, with physician input. At the bedside, physicians generally should act as patient advocates within constraints set by society, reasonable insurance coverage, and sound practice. Ad hoc rationing by the individual physician at the bedside may be inconsistent, discriminatory, and ineffective. In some cases, however, two patients may compete for the same limited resources, such as physician time or a bed in intensive care. When this occurs, physicians should nation their time and resources according to patients' medical needs and the probability of benefit.

ASSISTANCE WITH ETHICAL ISSUES. Discussing perplexing ethical issues with other members of the health care team, colleagues, or the hospital ethics committee often clarifies issues and suggests ways to improve communication and to deal with strong emotions. When struggling with difficult ethical issues, physicians may need to reevaluate their basic convictions, tolerate uncertainty, and maintain their integrity while respecting the opinions of others.

BIBLIOGRAPHY

- AJURIA A, LO B: When is ORT right? JAMA 273:156, 1995
- AMERICAN COLLEGE OF PHYSICIANS: American College of Physicians Ethics Manual. Ann Intern Med. 126:576, 1996
- BALCHER TL, CHALKS JR: Principles of Biomedical Ethics, 5th ed. New York, Oxford UP, 1997
- BALCHER TL, CHALKS JR: Principles of Biomedical Ethics, 5th ed. New York, Oxford UP, 1997
- CHALKS JR: Address to the prepared statements and ethics to physicians. JAMA 280:597, 1998
- CHALKS JR: Addressing Competence in Consent to Treatment: A Guide for Physicians and Other Health Professionals. New York, Oxford University Press, 1998
- KASSIRER JP: Managed care and the morality of the marketplace. N Engl J Med 333:50, 1995

reduces the likelihood of hyperthyroidism in a patient with paroxysmal atrial fibrillation.

While the representativeness and availability heuristics may play the major roles in shaping early diagnostic hypotheses, the acuity of a patient's illness can also be very influential. For example, clinicians are taught to consider aortic dissection routinely as a possible cause of acute severe chest discomfort along with myocardial infarction, even though the typical history of dissection is different from myocardial infarction and dissection is far less prevalent (Chap. 247). This recommendation is based on the recognition that a relatively rare but catastrophic diagnosis like aortic dissection is very difficult to make unless it is explicitly considered. If the clinician fails to elicit any of the characteristic features of dissection by history and finds equivalent blood pressures in both arms and no pulse deficits, he or she may feel comfortable in discarding the aortic dissection hypothesis. If, however, the chest x-ray shows a widened mediastinum, the hypothesis may be reinstated and a diagnostic test ordered [e.g., thoracic computed tomography (CT) scan, transesophageal echocardiogram] to evaluate it more fully. In noncritical situations, the prevalence of potential alternative diagnoses should play a much more prominent role in diagnostic hypothesis generation. The value of conducting a rapid systematic clinical survey of symptoms and organ systems to avoid missing important but inapparent clues cannot be overstated.

Because the generation and evaluation of appropriate diagnostic hypotheses is a skill that not all clinicians possess to an equal degree, errors in this process can occur, and in the patient with serious acute illness these may lead to tragic consequences. Consider the following hypothetical example. A 45-year-old male patient with a 3-week history of a "flu-like" upper respiratory infection (URI) presented to his physician with symptoms of dyspnea and a productive cough. Based on the presenting complaint, the clinician pulled out a "URI Assessment Form" to improve quality and efficiency of care. The physician quickly completed the examination components outlined on this structured form, noting in particular the absence of fever and a clear chest examination. He then prescribed an antibiotic for presumed bronchitis, showed the patient how to breathe into a paper bag to relieve his "hyperventilation," and sent him home with the reassurance that his illness was not serious. After a sleepless night with significant dyspnea unrelieved by rebreathing into a bag, the patient developed nausea and vomiting and collapsed. He was brought into the Emergency Department in cardiac arrest and could not be resuscitated. Autopsy showed a posterior wall myocardial infarction and a fresh thrombus in an atherosclerotic right coronary artery. What went wrong? The clinician decided, even before starting the history, that the patient's complaints were not serious. He therefore felt confident that he could perform an abbreviated and focused examination using the URI assessment protocol rather than considering the full range of possibilities and performing appropriate tests to confirm or refute his initial hypotheses. In particular, by concentrating on the "URI," the clinician failed to elicit the full dyspnea history, which would have suggested a far more serious disorder, and did not even search for other symptoms that could have directed him to the correct diagnosis.

This example illustrates how patients can diverge from textbook symptoms and the potential consequences of being unable to adapt the diagnostic process to real-world challenges. The expert, while recognizing that common things occur commonly, approaches each evaluation on high alert for clues that the initial diagnosis may be wrong. Patients often provide information that "does not fit" with any of the leading diagnostic hypotheses being considered. Distinguishing real clues from false trails can only be achieved by practice and experience. A less experienced clinician who tries to be too efficient (as in the above example) can make serious judgment errors.

MAJOR INFLUENCES ON CLINICAL DECISION-MAKING More than a decade of research on variations in clinician practice patterns has shed much light on forces that shape clinical

decisions. The use of heuristic "shortcuts," as detailed above, provides a partial explanation, but several other key factors play an important role in shaping diagnostic hypotheses and management decisions. These factors can be grouped conceptually into three overlapping categories: (1) factors related to physician personal characteristics and practice style, (2) factors related to the practice setting, and (3) economic incentive factors.

Practice Style Factors One of the key roles of the physician in medical care is to serve as the patient's agent to ensure that necessary care is provided at a high level of quality. Factors that influence this role include the physician's knowledge, training, and experience. It is obvious that physicians cannot practice evidence-based medicine if they are unfamiliar with the evidence. As would be expected, specialists generally know the evidence in their field better than do generalists. Surgeons may be more enthusiastic about recommending surgery than medical doctors because their belief in the beneficial effects of surgery is stronger. For the same reason, invasive cardiologists are much more likely to refer chest pain patients for diagnostic catheterization than are noninvasive cardiologists or generalists. The physician beliefs that drive these different practice styles are based on personal experience, recollection, and interpretation of the available medical evidence. For example, heart failure specialists are much more likely than generalists to achieve target angiotensin-converting enzyme (ACE) inhibitor therapy in their heart failure patients because they are more familiar with what the targets are (as defined by large clinical trials), have more familiarity with the specific drugs (including dosages and side effects), and are less likely to overreact to foreseeable problems in therapy such as a rise in creatinine levels or symptomatic hypotension. Other intriguing research has shown a wide distribution of acceptance times of antibiotic therapy for peptic ulcer disease following widespread dissemination of the "evidence" in the medical literature. Some gastroenterologists accepted this new therapy before the evidence was clear (reflecting, perhaps, an aggressive practice style), and some gastroenterologists lagged behind (a conservative practice style, associated in this case with older physicians). As a group, internists lagged several years behind gastroenterologists.

The opinion of influential leaders can also have an important effect on practice patterns. Such influence can occur at both the national level (e.g., expert physicians teaching at national meetings) and the local level (e.g., local educational programs, "curbside consultants"). Opinion leaders do not have to be physicians. When conducting rounds with clinical pharmacists, physicians are less likely to make medication errors and more likely to use target levels of evidence-based therapies.

The patient's welfare is not the only concern that drives clinical decisions. The physician's perception about the risk of a malpractice suit resulting from either an erroneous decision or a bad outcome creates a style of practice referred to as *defensive medicine*. This practice involves using tests and therapies with very small marginal returns to preclude future criticism in the event of an adverse outcome. For example, a 40-year-old woman who presents with a long-standing history of intermittent headache and a new severe headache along with a normal neurologic examination has a very low likelihood of structural intracranial pathology. Performance of a head CT or magnetic resonance imaging (MRI) scan in this situation would constitute defensive medicine. On the other hand, the results of the test could provide reassurance to an anxious patient.

Practice Setting Factors Factors in this category relate to the physical resources available to the physician's practice and the practice environment. *Physician-induced demand* is a term that refers to the repeated observation that physicians have a remarkable ability to accommodate to and employ the medical facilities available to them. A classic early study in this area showed that physicians in Boston had an almost 50% higher hospital admission rate than did physicians in New Haven, despite there being no obvious differences in the health of the cities' inhabitants. The physicians in New Haven were not aware of using fewer hospital beds for their patients, nor were the Boston physicians aware of using less stringent criteria to admit patients.

Table 3-1 Measures of Diagnostic Test Accuracy

Test Result	Disease Status	
	Present	Absent
Positive	True-positive (TP)	False-positive (FP)
Negative	False-negative (FN)	True-negative (TN)

IDENTIFICATION OF PATIENTS WITH DISEASE

True-positive rate (sensitivity) = $TP / (TP + FN)$
True-negative rate = $TN / (TN + FP)$
True-positive rate = 1 - false-negative rate

IDENTIFICATION OF PATIENTS WITHOUT DISEASE

True-negative rate (specificity) = $TN / (TN + FP)$
True-positive rate = $FP / (TN + FP)$
True-negative rate = 1 - false-positive rate

(1 - specificity). A perfect test would have a sensitivity of 100% and a specificity of 100% and would completely separate patients with disease from those without it.

Calculating sensitivity and specificity require selection of a cut-point value for the test to separate "normal" from "diseased" subjects. As the cutpoint is moved to improve sensitivity, specificity typically falls and vice versa. This dynamic tradeoff between more accurate identification of subjects with versus those without disease is often displayed graphically as a receiver operating characteristic (ROC) curve. An ROC curve plots sensitivity (y-axis) versus 1 - specificity (x-axis). Each point on the curve represents a potential cutpoint with an associated sensitivity and specificity value. The area under the ROC curve is often used as a quantitative measure of the information content of a test. Values range from 0.5 (no diagnostic information at all, test is equivalent to flipping a coin) to 1.0 (perfect test).

In the diagnostic testing literature, ROC areas are often used to compare alternative tests. The test with the highest area (i.e., closest to 1.0) is presumed to be the most accurate. However, ROC curves are not a panacea for evaluation of diagnostic test utility. Like Bayes' theorem, they are typically focused on only one possible test parameter (e.g., ST segment response in a treadmill exercise test) to the exclusion of other potentially relevant data. In addition, ROC area comparisons do not simulate the way test information is actually used in clinical practice. Finally, biases in the underlying population used to generate the ROC curves (e.g., related to an unrepresentative test sample) can bias the ROC area and the validity of a comparison among tests.

Measures of Disease Probability and Bayes' Theorem. Unfortunately, there are no perfect tests; after every test is completed the true disease state of the patient remains uncertain. Quantifying this residual uncertainty can be done with Bayes' theorem. This theorem provides a simple mathematical way to calculate the posterior probability of disease from three parameters: the pretest probability of disease, the test sensitivity, and the test specificity (Table 3-2). The pretest probability is a quantitative expression of the confidence in a diagnosis before the test is performed. In the absence of more relevant information it is usually estimated from the prevalence of the disease in the underlying population. For some common conditions, such as coronary artery disease (CAD), nomograms and statistical models have been created to generate better estimates of pretest probability from elements of the history and physical examination. The posterior probability, then, is a revised statement of the confidence in the diagnosis, taking into account both what was known before and after the test.

To understand how Bayes' theorem creates this revised confidence statement, it is useful to examine a nomogram version of Bayes' theorem that uses the same three parameters to predict the posttest probability of disease (Fig. 3-1). In this nomogram, the accuracy of the diagnostic test in question is summarized by the likelihood ratio for a

Other environmental factors that can influence decision-making include the local availability of specialists for consultations and procedures, "high tech" facilities such as angiography suites, a heart surgery program, and MRI machines.

Economic Incentives. Economic incentives are closely related to the other two categories of practice-modifying factors. Financial issues can exert both stimulatory and inhibitory influences on clinical practice. In general, physicians are paid on a fee-for-service capitation, or salary basis (Chap. 4). In fee-for-service, the more the physician does, the more the physician gets paid. The incentive in this case is to do more. When fees are reduced (discounted fee-for-service), doctors tend to increase the number of services billed for. Capitation, in contrast, provides a fixed payment per patient per year, encouraging physicians to take on more patients but to provide each patient with fewer services. Expensive services are more likely to be affected by this type of incentive than inexpensive preventive services. Salary components of incentive plans vary, sometimes the same regardless of the amount of clinical work performed. The incentive here is to see fewer patients. Recognizing these powerful shapes of physician behavior, managed care plans have begun to explore combinations of the three reimbursement types with the goal of improving individual physician productivity while restraining their use of expensive tests and drugs.

In summary, expert clinical decision-making can be appreciated as a complex interplay between cognitive devices used to simplify large amounts of complex information interacting with physician biases regarding education, training, and experience, all of which are shaped by powerful, sometimes perverse, external forces. In the next section, we will review a set of statistical tools and concepts that can assist in making clinical decisions under uncertainty.

QUANTITATIVE METHODS TO AID CLINICAL DECISION-MAKING

The process of medical decision-making can be divided into two parts: (1) defining the available courses of action and estimating the likely outcomes with each, and (2) assessing the desirability of the outcomes. The former task involves integrating key information about the patient along with relevant evidence from the medical literature to create the likelihood of a decision problem. The remainder of this chapter will present some quantitative tools to assist the clinician in these activities. These tools can be divided into those that assist the clinician in making better outcome predictions, which are then used to make decisions, and those that support the decision process directly. While these tools are not yet used routinely in daily clinical practice, the computerization of medicine is creating the required substrate for their future widespread dissemination.

QUANTITATIVE MEDICAL PREDICTIONS. Diagnostic Testing. The purpose of performing a test on a patient is to reduce uncertainty about the patient's diagnosis or prognosis and to aid the clinician in making management decisions. Although diagnostic tests are commonly thought of as laboratory tests (e.g., measurement of serum amylase level) or procedures (e.g., colonoscopy or bronchoscopy), any technology that changes our understanding of the patient's problem qualifies as a diagnostic test. Thus, even the history and physical examination can be considered a form of diagnostic test. In clinical medicine, it is common to reduce the results of a test to a dichotomous outcome, such as positive or negative, normal or abnormal. In many cases, such simplification results in the waste of useful information. However, such simplification makes it easier to demonstrate some of the quantitative ways in which test data can be used.

To characterize the accuracy of diagnostic tests, four terms are commonly used (Table 3-1). The true-positive rate, i.e., the sensitivity, provides a measure of how well the test correctly identifies patients with disease. The false-negative rate is calculated as $(1 - \text{sensitivity})$. The true-negative rate, i.e., the specificity, reflects how well the test correctly identifies patients without disease. The false-positive rate is

decisions. The use of heuristic "shortcuts," as detailed above, provides a partial explanation, but several other key factors play an important role in shaping diagnostic hypotheses and management decisions. These factors can be grouped conceptually into three overlapping categories: (1) factors related to physician personal characteristics and practice style, (2) factors related to the practice setting, and (3) economic incentive factors.

Practice Style Factors. One of the key roles of the physician in medical care is to serve as the patient's agent to ensure that necessary care is provided at a high level of quality. Factors that influence this role include the physician's knowledge, training, and experience. It is obvious that physicians cannot practice evidence-based medicine if they are unfamiliar with the evidence. As would be expected, specialists generally know the evidence in their field better than do generalists. Surgeons may be more enthusiastic about recommending surgery than medical doctors because their belief in the beneficial effects of surgery is stronger. For the same reason, invasive cardiologists are much more likely to refer chest pain patients for diagnostic catheterization than are noninvasive cardiologists or generalists. The physician beliefs that drive these different practice styles are based on personal experience, recollection, and interpretation of the available medical evidence. For example, heart failure specialists are much more likely than generalists to achieve target angiotensin-converting enzyme (ACE) inhibitor therapy in their heart failure patients because they are more familiar with what the targets are (as defined by large clinical trials), have more familiarity with the specific drugs (including dosages and side effects), and are less likely to overreact to foreseeable problems in therapy such as a rise in creatinine levels or symptomatic hypotension. Other intriguing research has shown a wide distribution of acceptance times of antibiotic therapy for peptic ulcer disease following widespread dissemination of the "evidence" in the medical literature. Some gastroenterologists accepted this new therapy before the evidence was clear (reflecting, perhaps, an aggressive practice style), and some gastroenterologists lagged behind (a conservative practice style), associated in this case with older physicians). As a group, internists lagged several years behind gastroenterologists.

The opinion of influential leaders can also have an important effect on practice patterns. Such influence can occur at both the national level (e.g., expert physicians teaching at national meetings) and the local level (e.g., local educational programs, "outreach consultants"). Opinion leaders do not have to be physicians. When conducting rounds with clinical pharmacists, physicians are less likely to make medication errors and more likely to use target levels of evidence-based therapies. The patient's welfare is not the only concern that drives clinical decisions. The physician's perception about the risk of a malpractice suit resulting from either an erroneous decision or a bad outcome creates a style of practice referred to as defensive medicine. This practice involves using tests and therapies with very small marginal returns to preclude future criticism in the event of an adverse outcome. For example, a 40-year-old woman who presents with a long-standing history of intermittent headache and a new severe headache along with a normal neurological examination has a very low likelihood of structural intracranial pathology. Performance of a head CT or magnetic resonance imaging (MRI) scan in this situation would constitute defensive medicine. On the other hand, the results of the test could provide reassurance to an anxious patient.

Practice Setting Factors. Factors in this category relate to the physical resources available to the physician's practice and the practice environment. Physician-induced demand is a term that refers to the repeated observation that physicians have a remarkable ability to accommodate to and employ the medical facilities available to them. A classic early study in this area showed that physicians in Boston had an almost 30% higher hospital admission rate than did physicians in the cities' inhabitants. The physicians in New Haven were not aware of using fewer hospital beds for their patients, nor were the Boston physicians aware of using less stringent criteria to admit patients.

10
reduces the likelihood of hyperthyroidism in a patient with paroxysmal atrial fibrillation.

While the representativeness and availability hypotheses may play the major roles in shaping early diagnostic hypotheses, the acuity of a patient's illness can also be very influential. For example, clinicians are taught to consider aortic dissection routinely as a possible cause of acute severe chest discomfort along with myocardial infarction, even though the typical history of dissection is different from myocardial infarction and dissection is far less prevalent (Chap. 247). This recommendation is based on the recognition that a relatively rare but catastrophic diagnosis like aortic dissection is very difficult to make unless it is explicitly considered. If the clinician fails to elicit any of the characteristic features of dissection by history and finds equivalent blood pressures in both arms and no pulse deficits, he or she may feel comfortable in discarding the aortic dissection hypothesis. If, however, the chest x-ray shows a widened mediastinum, the hypothesis may be reinstated and a diagnostic test ordered (e.g., thoracic computed tomography [CT] scan, transesophageal echocardiogram) to evaluate it more fully. In noncritical situations, the prevalence of potential alternative diagnoses should play a much more prominent role in diagnostic hypothesis generation. The value of conducting a rapid systematic clinical survey of symptoms and organ systems to avoid missing important but apparent clues cannot be overstated.

Because the generation and evaluation of appropriate diagnostic hypotheses is a skill that not all clinicians possess to an equal degree, errors in this process can occur, and in the patient with serious acute illness these may lead to tragic consequences. Consider the following hypothetical example. A 45-year-old male patient with a 3-week history of a "hilar," upper respiratory infection (URI) presented to his physician with symptoms of dyspnea and a productive cough. Based on the presenting complaint, the clinician pulled out a "URI Assessment Form" to improve quality and efficiency of care. The physician quickly completed the examination components outlined on this structured form, noting in particular the absence of fever and a clear chest examination. He then prescribed an antibiotic for presumed bronchitis, showed the patient how to breathe into a paper bag to relieve his "hyperventilation," and sent him home with the reassurance that his illness was not serious. After a sleepless night with significant dyspnea and unrelieved by rebreathing into a bag, the patient developed nausea and vomiting and collapsed. He was brought into the Emergency Department in cardiac arrest and could not be resuscitated. Autopsy showed a posterior wall myocardial infarction and a fresh thrombus in an atherosclerotic right coronary artery. What went wrong? The clinician decided, even before starting the history, that the patient's complaints were not serious. He therefore felt confident that he could perform an abbreviated and focused examination using the URI assessment protocol rather than considering the full range of possibilities and performing appropriate tests to confirm or refute his initial hypotheses. In particular, by concentrating on the "URI," the clinician failed to elicit the full dyspnea history, which would have suggested a far more serious disorder, and did not even search for other symptoms that could have directed him to the correct diagnosis.

This example illustrates how patients can diverge from textbook symptoms and the potential consequences of being unable to adapt the diagnostic process to real-world challenges. The expert, while recognizing that common things occur commonly, approaches each evaluation on high alert for clues that the initial diagnosis may be wrong. Patients often provide information that "does not fit" with any of the leading diagnostic hypotheses being considered. Distinguishing real clues from false trails can only be achieved by practice and experience. A less experienced clinician who tries to be too efficient (as in the above example) can make serious judgment errors.

MAJOR INFLUENCES ON CLINICAL DECISION-MAKING. More than a decade of research on variations in clinician practice patterns has shed much light on forces that shape clinical

Table 3-2 Measures of Disease Probability

Pretest probability of disease = probability of disease before test is done; may use population prevalence of disease or more patient-specific data to generate this probability estimate.

Posttest probability of disease = probability of disease accounting for both pretest probability and test results; also called predictive value of the test.

Bayes' theorem

Computational version:

$$\text{Posttest probability} = \frac{\text{Pretest probability} \times \text{test sensitivity}}{\text{Pretest probability} \times \text{test sensitivity} + (1 - \text{disease prevalence}) \times \text{test false-positive rate}}$$

Example [with a pretest probability of 0.50 and a "positive" diagnostic test result (test sensitivity = 0.90, test specificity = 0.90)]:

$$\begin{aligned} \text{Posttest probability} &= \frac{(0.50)(0.90)}{(0.50)(0.90) + (0.50)(0.10)} \\ &= 0.90 \end{aligned}$$

positive test, which is the ratio of the true-positive rate to the false-positive rate [or sensitivity/(1 - specificity)]. For example, a test with a sensitivity of 0.90 and a specificity of 0.90 has a likelihood ratio of 0.90/(1 - 0.90), or 9. Thus, for this hypothetical test, a "positive"




FIGURE 3-1 Nomogram version of Bayes' theorem used to predict the posttest probability of disease (left-hand scale) using the pretest probability of disease (right-hand scale) and the likelihood ratio for a positive test (middle scale). The likelihood ratio is calculated as the sensitivity/(1 - specificity). To use, place a straight edge connecting the pretest probability and the likelihood ratio, and read off the posttest probability. This figure illustrates the value of a positive exercise treadmill test (likelihood ratio 4) and a positive exercise thallium SPECT study (likelihood ratio 9) in the patient with a pretest probability of coronary artery disease of 50%. Treadmill results shown in solid line; thallium results in dashed line. (Adapted from Fagan TJ: *N Engl J Med* 293:257, 1975.)

result is 9 times more likely in a patient with the disease than in a patient without it. The more accurate the test, the higher the likelihood ratio. However, if sensitivity is excellent but specificity is less so, the likelihood ratio will be substantially reduced (e.g., with a 90% sensitivity but a 60% specificity, the likelihood ratio is 2.25). Most tests in medicine have likelihood ratios for a positive result between 1.5 and 20.

Consider two tests commonly used in the diagnosis of CAD, an exercise treadmill and an exercise thallium-201 single photon emission CT (SPECT) test (Chap. 244). Meta-analysis has shown the treadmill to have an average sensitivity of 66% and an average specificity of 84%, yielding a likelihood ratio of 4.1 [0.66/(1 - 0.84)]. If we use this test on a patient with a pretest probability of CAD of 10%, the posttest probability of disease following a positive result rises only to about 30%. If a patient with a pretest probability of CAD of 80% has a positive test result, the posttest probability of disease is about 95%.

The exercise thallium SPECT test is a more accurate test for the diagnosis of CAD. For our purposes, assume that it has both a sensitivity and specificity of 90%, yielding a likelihood ratio of 9.0 [0.90/(1 - 0.90)]. If we again test our low pretest probability patient and he has a positive test, using Fig. 3-1 we can demonstrate that the posttest probability of CAD rises from 10 to 50%. However, from a decision-making point of view, the more accurate test has not been able to improve diagnostic confidence enough to change management. In fact, the test has moved us from being fairly certain that the patient did not have CAD to being completely undecided (a 50:50 chance of disease). In a patient with a pretest probability of 80%, using the more accurate thallium SPECT test raises the posttest probability to 97% (compared with 95% for the exercise treadmill). Again, the more accurate test does not provide enough improvement in posttest confidence to alter management, and neither test has improved much upon what was known from clinical data alone.

If the pretest probability is low (e.g., $\leq 20\%$), even a positive result on a very accurate test will not move the posttest probability to a range high enough to rule in disease (e.g., $\geq 80\%$). Conversely, with a high pretest probability, a negative test will not adequately rule out disease. Thus, the largest gain in diagnostic confidence from a test occurs when the clinician is most uncertain before performing it (e.g., pretest probability between 30 and 70%). For example, if a patient has a pretest probability for CAD of 50%, a positive exercise treadmill test will move the posttest probability to 80% and a positive exercise thallium SPECT test will move it to 90% (Fig. 3-1).

Bayes' theorem, as presented above, employs a number of important simplifications that should be considered. First, few tests have only two useful outcomes, positive or negative, and many tests provide numerous pieces of data about the patient. Even if these can be integrated into a summary result, multiple levels of useful information may be present (e.g., strongly positive, positive, indeterminate, negative, strongly negative). While Bayes' theorem can be adapted to this more detailed test result format, it is computationally complex to do so. Second, Bayes' theorem assumes that the information from the test is completely unique and nonoverlapping with information used to estimate the pretest probability. This independence assumption, however, is often wrong. In many cases, test results are correlated with patient characteristics. For example, the findings of cardiomegaly and pulmonary edema on chest x-ray are correlated with the historic features of heart failure and with the physical findings of a displaced left ventricular apical impulse, an S₃ gallop, and rales. The unique predictive information contributed by the test in this case (the chest x-ray) is only a fraction of its total information because much had already been learned about the probability of heart failure before the test was done.

Finally, it has long been thought that sensitivity and specificity are prevalence-independent parameters of test accuracy, and many texts still make this assertion. This statistically useful assumption, however, is clinically wrong. For example, a treadmill exercise test has a sensitivity in a population of patients with one-vessel CAD of around 30%, whereas the sensitivity in severe three-vessel CAD approaches

Table 3-2 Measures of Disease Probability

Pretest probability of disease = probability of disease before test is done; may use population prevalence of disease or more patient-specific data to generate this probability estimate.

Posttest probability of disease = probability of disease accounting for both pretest probability and test results; also called predictive value of the test.

Bayes' theorem

$$\text{Posttest probability} = \frac{\text{Pretest probability} \times \text{test sensitivity}}{(\text{Pretest probability} \times \text{test sensitivity}) + (1 - \text{Pretest probability}) \times \text{test false-positive rate}}$$

Example (with a pretest probability of 0.90 and a "positive" diagnostic test result [test sensitivity = 0.90, test specificity = 0.90]):

$$\text{Posttest probability} = \frac{(0.90)(0.90)}{(0.90)(0.90) + (0.10)(0.10)} = 0.90$$

positive test, which is the ratio of the true-positive rate to the false-positive rate (or sensitivity/(1 - specificity)). For example, a test with a sensitivity of 0.90 and a specificity of 0.90 has a likelihood ratio of 0.90/(1 - 0.90), or 9. Thus, for this hypothetical test, a "positive"




FIGURE 3-1 Nomogram version of Bayes' theorem used to predict the posttest probability of disease (left-hand scale) using the pretest probability of disease (right-hand scale) and the likelihood ratio for a positive test (middle scale). The likelihood ratio is calculated as the sensitivity/(1 - specificity). To use, place a straight edge connecting the pretest probability and the likelihood ratio, and read off the posttest probability. This figure illustrates the value of a positive exercise treadmill test (likelihood ratio 9) and a positive exercise thallium SPECT study (likelihood ratio 9) in the patient with a pretest probability of coronary artery disease of 50%. Treadmill results shown in solid line; thallium results in dashed line. (Adapted from Fagan *ET: N Engl J Med* 331:257, 1975.)

result is 9 times more likely in a patient with the disease than in a patient without it. The more accurate the test, the higher the likelihood ratio. However, if sensitivity is excellent but specificity is less so, the likelihood ratio will be substantially reduced (e.g., with a 90% sensitivity but a 60% specificity, the likelihood ratio is 2.25). Most tests in medicine have likelihood ratios for a positive result between 1.5 and 20.

Consider two tests commonly used in the diagnosis of CAD, an exercise treadmill and an exercise thallium-201 single-photon emission CT (SPECT) test (Chap. 244). Meta-analysis has shown the treadmill to have an average sensitivity of 66% and an average specificity of 84%, yielding a likelihood ratio of 4.1 [0.66/(1 - 0.84)]. If we use this test on a patient with a pretest probability of CAD of 10%, the posttest probability of disease following a positive result rises only to about 30%. If a patient with a pretest probability of disease is about 95%, a positive test result, the posttest probability of disease is about 95%.

The exercise thallium SPECT test is a more accurate test for the diagnosis of CAD. For our purposes, assume that it has both a sensitivity and specificity of 90%, yielding a likelihood ratio of 9.0 [0.90/(1 - 0.90)]. If we again test our low pretest probability patient and he has a positive test result, using Fig. 3-1 we can demonstrate that the posttest probability of CAD rises from 10 to 50%. However, from a decision-making point of view, the more accurate test has not been able to improve diagnostic confidence enough to change management. In fact, the test has moved us from being fairly certain that the patient did not have CAD to being completely undecided (a 50-50 chance of disease). In a patient with a pretest probability of 80%, using the more accurate thallium SPECT test raises the posttest probability to 97% (compared with 95% for the exercise treadmill). Again, the more accurate test does not provide enough improvement in posttest confidence to alter management, and neither test has improved much upon what was known from clinical data alone.

If the pretest probability is low (e.g., $\leq 20\%$), even a positive result on a very accurate test will not move the posttest probability to a range high enough to rule in disease (e.g., $\geq 80\%$). Conversely, with a high pretest probability, a negative test will not adequately rule out disease. Thus, the largest gain in diagnostic confidence from a test occurs when the clinician is most uncertain before performing it (e.g., pretest probability between 30 and 70%). For example, if a patient has a pretest probability for CAD of 50%, a positive exercise treadmill test will move the posttest probability to 80% and a positive exercise thallium SPECT test will move it to 90% (Fig. 3-1).

Bayes' theorem, as presented above, employs a number of important simplifications that should be considered. First, few tests have only two useful outcomes, positive or negative, and many tests provide numerous pieces of data about the patient. Even if these can be integrated into a summary result, multiple levels of useful information may be present (e.g., strongly positive, positive, indeterminate, negative, strongly negative). While Bayes' theorem can be adapted to this more detailed test result format, it is computationally complex to do so. Second, Bayes' theorem assumes that the information from the test is completely unique and nonoverlapping with information used to estimate the pretest probability. This independence assumption, however, is often wrong. In many cases, test results are correlated with patient characteristics. For example, the findings of cardiomegaly and pulmonary edema on chest x-ray are correlated with the historic features of heart failure and with the physical findings of a displaced left ventricular apical impulse, an S₃ gallop, and rales. The unique predictive information contributed by the test in this case (the chest x-ray) is only a fraction of its total information because much had already been learned about the probability of heart failure before the test was done.

Finally, it has long been thought that sensitivity and specificity are prevalence-independent parameters of test accuracy, and many texts still make this assertion. This statistically useful assumption, however, is clinically wrong. For example, a treadmill exercise test has a sensitivity in a population of patients with one-vessel CAD of around 30%, whereas the sensitivity in severe three-vessel CAD approaches

90%. Thus, the best estimate of sensitivity to use in a particular decision will often vary depending on the distribution of disease stages present in the patient population. A hospitalized population typically has a higher prevalence of disease and in particular a higher prevalence of more advanced disease stages than an outpatient population. As a consequence, test sensitivity will tend to be higher in hospitalized patients, whereas test specificity will be higher in outpatients.

Statistical Prediction Models Bayes' theorem, as presented above, deals with a clinical prediction problem that is unrealistically simple relative to most problems a clinician faces. Prediction models based on multivariable statistical models can handle much more complex problems and substantially enhance predictive accuracy for specific situations. Their particular advantage is the ability to take into account many overlapping pieces of information and assign a relative weight to each based on its unique contribution to the prediction in question. For example, a logistic regression model to predict the probability of CAD takes into account all of the relevant independent factors from the clinical examination and diagnostic testing in a way that is more complex than the use of Bayes' theorem. However, despite this strength, the models are too complex computationally to use without a calculator or computer (although this limitation may be overcome when medicine is practiced from a fully computerized platform). To date, only a handful of prediction models have been developed and properly validated. The importance of independent validation in a population separate from the one used to develop the model cannot be overstated. Unfortunately, most published models have not been properly validated, making their utility in clinical practice questionable at best.

When statistical models have been compared directly with expert clinicians, they have been found to be more consistent, as would be expected, but not significantly more accurate. Their biggest promise, then, would seem to be to make less-experienced clinicians more accurate predictors of outcome.

DECISION SUPPORT TOOLS

DECISION SUPPORT SYSTEMS: Over the past 30 years, many attempts have been made to develop computer systems to help clinicians make decisions and manage patients. Conceptually, computers offer a very attractive way to handle the vast information load of today's physicians. The computer can help by making accurate predictions of outcome, simulating the whole decision process, or providing statistical guidance. Computer-based prediction using Bayesian or statistical regression models inform a clinical decision but do not usually reach a "conclusion" or "recommendation." Artificial intelligence systems attempt to simulate or replace human reasoning with a "computer-based" analogue. To date, such approaches have achieved only limited success. Reminder or protocol-directed systems do not make predictions but use existing algorithms, such as practice guidelines, to guide clinical practice. In general, however, decision




FIGURE 3-2 Decision model used to evaluate strategies for management of the risk of infective endocarditis after catheter-associated *Staphylococcus aureus* bacteremia. The square node indicates a decision between possible management strategies, round nodes represent chance events, and rectangular (or terminal) nodes indicate the outcomes of interest. All nonterminal chance nodes in the main tree (structure A) enter substructure B. All nonterminal chance nodes in substructure B enter substructure C. TEE, transthoracic echocardiography. (From Rosen *et al*.)

support systems have shown little impact on practice. Reminder systems, although not yet in widespread use, have shown the most promise, particularly in correcting drug dosing and in promoting guideline adherence. The full potential of these approaches will only be achieved when computers are fully integrated into medical practice.

DECISION ANALYSIS Compared with the methods discussed above, decision analysis represents a completely different approach to decision support. Its principal application is in decision problems that are complex and involve a substantial risk, a high degree of uncertainty in some key area, or an idiosyncratic feature that does not "fit" the available evidence. Three general steps are involved. First, the decision problem must be clearly defined. Second, the elements of the decision must be made explicit. This involves specifying the alternatives being considered, their relevant outcomes, the probabilities attached to each outcome, and the relative desirability (called "utility") of each outcome. Cost can also be assigned to each branch of the decision tree, allowing calculation of cost-effectiveness (Chap. 4).

An example of a decision tree used to evaluate strategies for management of the risk of infective endocarditis after catheter-associated *Staphylococcus aureus* bacteremia is shown in Fig. 3-2. Approximately 35,000 cases of *S. aureus* bacteremia occur each year in the United States. The development of complicating endocarditis, which occurs in about 6% of cases, is associated with high mortality (31% mortality, 21% stroke rate) and morbidity. The three choices for management of the bacteremia are (1) transthoracic echocardiography (TEE), (2) a 4-week course of intravenous antibiotics (long-course), or (3) a 2-week course of intravenous antibiotics (short-course). In the TEE strategy, a 4-week course is given if it is not. With

each strategy, there is a risk that the patient will develop endocarditis with or without major complications. In this analysis, the longest quality-adjusted survival (5.47 quality-adjusted life-years) was associated with the 4-week antibiotic course strategy, which also had the highest costs (\$14,136 per patient), whereas the lowest costs (\$9830 per patient) and worst outcomes (5.42 quality-adjusted life-years) were associated with the 2-week antibiotic course strategy. From a clinical point of view (ignoring costs), the 4-week antibiotic course was best. From a cost-effectiveness point of view, the TEE strategy (5.46 quality-adjusted life-years and \$10,051 per patient costs) provided the best balance of added benefits and costs. Thus, decision analysis can be extremely helpful in clarifying tradeoffs in outcomes and costs in difficult management areas such as the above where it is highly unlikely that an adequate randomized trial will ever be done.

The data needed to fill in a decision tree (Fig. 3-2) are typically cobbled together from a variety of sources, including the literature (randomized trials, meta-analyses, observational studies) and expert opinion. Once the decision tree is finished, the decision is "analyzed" by calculating the average value of each limb of the tree. The decision arm with the highest net value (or expected utility) is the preferred choice. The value of this exercise, however, is not so much in developing a prescription for action as it is in exploring the key elements and pressure points of a complex or difficult decision. The process of building the decision tree forces the analyst to be explicit about the choices being considered and all their relevant outcomes. Areas of high uncertainty are readily identified. Sensitivity analyses are an integral part of decision analysis and involve systematically varying the value of each key parameter in the model alone (one-way sensitivity analysis), in pairs (two-way), or in higher combinations (multivariable) to assess the impact on choice of preferred management strategy. In the above example, varying the incidence of endocarditis resulting from *S. aureus* bacteremia from 3% to over 50% had no impact on the choice of TEE as the preferred strategy.

User friendly personal computer-based software packages now make the creation and analysis of decision trees much more straightforward than in the past. However, the process is still too cumbersome and time-consuming to be used on a routine basis. When medicine is practiced from a fully computerized platform, a library of prestructured decision trees with user modifiable values can be made available to support practitioners working with individual patients.

CONCLUSIONS

In this era of evidence-based medicine, it is tempting to think that all the difficult decisions practitioners face have been or soon will be solved and digested into practice guidelines and computerized reminders. For the foreseeable future, however, such is not the case. Meta-analyses cannot generate evidence where there are no adequate randomized trials, and most of what clinicians face will never be thoroughly tested in a randomized trial. Excellent clinical reasoning skills and experience supplemented by well-designed quantitative tools and a keen appreciation for individual patient preferences will continue to be of paramount importance in the professional life of medical practitioners for years to come.

BIBLIOGRAPHY

- AYANIAN JZ et al: Knowledge and practices of generalist and specialist physicians regarding drug therapy for acute myocardial infarction. *N Engl J Med* 331:1136, 1994
 EDDY DM: Anatomy of a decision. *JAMA* 263:441, 1990
 HIRTH RA et al: Specialist and generalist physicians' adoption of antibiotic therapy to eradicate *Helicobacter pylori* infection. *Med Care* 34:1199, 1996
 KASSIRER JP, KOPELMAN RI: *Learning Clinical Reasoning*. Baltimore, Williams & Wilkins, 1991
 NAYLOR CD: Gray zones of clinical practice: Some limits to evidence-based medicine. *Lancet* 345:840, 1995

- ROSEN AB et al: Cost-effectiveness of transesophageal echocardiography to determine the duration of therapy for intravascular catheter-associated *Staphylococcus aureus* bacteremia. *Ann Intern Med* 130:810, 1999
 SCHULMAN KA et al: The effect of race and sex on physicians' recommendations for cardiac catheterization. *N Engl J Med* 340:618, 1999
 SEKKARJE MA, MOSS AH: Withholding and withdrawing dialysis: The role of physician specialty and education and patient functional status. *Am J Kidney Dis* 31:464, 1998

4

Daniel B. Mark

ECONOMIC ISSUES IN CLINICAL MEDICINE

The United States has the distinction of having some of the best medical care of any technologically advanced country. We have many of the best hospitals and doctors in the world. The research pipeline is full of significant new therapeutic advances, with revolutionary genetic-based therapies perhaps only a decade away. Our citizens largely subscribe to the principle that excellent medical care should be available to all, regardless of ability to pay. Yet we also have over 43 million people (most of them employed and earning minimal wages) without any health insurance and many more who are inadequately insured. Since the collapse of the Clinton health care reform efforts in 1994, U.S. health policy has been directed by marketplace forces that have created powerful and sometimes perverse incentives in medicine: Health insurance companies that use every available means to avoid insuring sick people; "managed care" programs that really only manage costs; doctors who are provided incentives to provide less medical care; and pharmaceutical companies that develop powerful and expensive new drugs priced beyond the reach of many of the elderly and chronically ill who need them most.

Facing such powerful and chaotic forces, physicians tend to focus narrowly on what they are most comfortable with, taking care of individual patients and conducting academic investigations. Many doctors consider economics too arcane for them to grasp and therefore do not even try. Consequently, when presented with economic arguments and evidence they are often unable to discriminate the legitimate from the fallacious. More importantly, they are ill equipped to defend their patients' interests in the crucible of cost containment that characterizes the modern managed care era.

This chapter has two goals: first, to provide a brief introduction to some of the larger economic forces that shape modern medical practices, and second, to introduce the economic tools that are used for assessing the value of medical practices, including cost effectiveness analysis.

HEALTH CARE SPENDING AND FINANCING

HOW MUCH IS SPENT ON HEALTH CARE? In 1997, the United States spent \$1.1 trillion on its health care system, representing 13.5% of the gross domestic product (GDP) (a crude measure of national income). Most of this (\$969 billion) was spent on personal health care: 34% went to hospitals, 20% to physicians, 7% to nursing homes, and 8% to outpatient pharmaceuticals. In comparison, Canada and Western European countries spend a substantially smaller portion (6 to 10%) of their national income on health care but their citizens appear to be equally healthy, at least by crude metrics such as life expectancy and infant mortality rates. Economists and politicians have for years used such data to argue that the United States spends too much on health care. The issue of how much to spend is an inherently political one, however, and the discipline of economics has little to say about it.

WHO PAYS FOR HEALTH CARE? Two major factors are continually driving up the costs of medical care: introduction into medical practice of new medical technologies (drugs, devices, procedures) that have a high price tag, and the aging of the U.S. population

sincere concern, the willingness to take the time to explain all aspects of the illness, and a nonjudgmental attitude when dealing with patients whose cultures, lifestyles, attitudes, and values differ from those of the physician are just a few of the characteristics of the humane physician. Every physician will, at times, be challenged by patients who evoke strongly negative (or strongly positive) emotional responses. Physicians should be alert to their own reactions to such patients and situations and should consciously monitor and control their behavior so that the patients' best interests remain the principal motivation for their actions at all times.

An important aspect of patient care involves an appreciation of the "quality of life," a subjective assessment of what each patient values most. Such an assessment requires detailed, sometimes intimate knowledge of the patient, which can usually be obtained only through deliberate, unhurried, and often repeated conversations. It is in these situations that the time constraints of a managed care setting may prove problematic.

The famous statement of Dr. Francis Peabody is even more relevant today than when delivered more than three-quarters of a century ago:

The significance of the intimate personal relationship between physician and patient cannot be too strongly emphasized, for in an extraordinarily large number of cases both the diagnosis and treatment are directly dependent on it. One of the essential qualities of the clinician is interest in humanity, for the secret of the care of the patient is in caring for the patient.

CLINICAL SKILLS History Taking The written history of an illness should embody all the facts of medical significance in the life of the patient. Recent events should be given the most attention. The patient should, at some point, have the opportunity to tell his or her own story of the illness without frequent interruption and, when appropriate, receive expressions of interest, encouragement, and empathy from the physician. The physician must be alert to the possibility that any event related by the patient, however trivial or apparently remote, may be the key to the solution of the medical problem.

An informative history is more than an orderly listing of symptoms; something is always gained by listening to patients and noting the way in which they describe their symptoms. Inflections of voice, facial expression, gestures, and attitude may reveal important clues to the meaning of the symptoms to the patient. Taking history often involves much data gathering. Patients vary in their medical sophistication and ability to recall facts. Medical history should therefore be corroborated whenever possible. The family and social history can also provide important insights into the types of diseases that should be considered. In listening to the history, the physician discovers not only something about the disease but also something about the patient. The process of history taking provides an opportunity to observe the patient's behavior and to watch for features to be pursued more thoroughly during the physical examination.

The very act of eliciting the history provides the physician with the opportunity to establish or enhance the unique bond that is the basis for the ideal patient-physician relationship. It is helpful to develop an appreciation of the patient's perception of the illness, the patient's expectations of the physician and the medical care system, and the financial and social implications of the illness to the patient. The confidentiality of the patient-physician relationship should be emphasized, and the patient should be given the opportunity to identify any aspects of the history that should not be disclosed.

Physical Examination Physical signs are objective indications of disease whose significance is enhanced when they confirm a functional or structural change already suggested by the patient's history. At times, however, the physical signs may be the only evidence of disease.

The physical examination should be performed methodically and thoroughly, with consideration for the patient's comfort and modesty. Although attention is often directed by the history to the diseased organ

or part of the body, the examination of a new patient must extend from head to toe in an objective search for abnormalities. Unless the physical examination is systematic, important segments may be omitted. The results of the examination, like the details of the history, should be recorded at the time they are elicited, not hours later when they are subject to the distortions of memory. Skill in physical diagnosis is acquired with experience, but it is not merely technique that determines success in eliciting signs. The detection of a few scattered petechiae, a faint diastolic murmur, or a small mass in the abdomen is not a question of keener eyes and ears or more sensitive fingers but of a mind alert to these findings. Since physical findings are subject to changes, the physical examination should be repeated as frequently as the clinical situation warrants.

Laboratory Tests The availability of a wide array of laboratory tests has increased our reliance on these studies for the solution of clinical problems. The accumulation of laboratory data does not relieve the physician from the responsibility of careful observation, examination, and study of the patient. It is also essential to bear in mind the limitations of such tests. By virtue of their impersonal quality, complexity, and apparent precision, they often gain an aura of authority regardless of the fallibility of the tests themselves, the instruments used in the tests, and the individuals performing or interpreting them. Physicians must weigh the expense involved in the laboratory procedures they order relative to the value of the information they are likely to provide.

Single laboratory tests are rarely ordered. Rather, they are generally obtained as "batteries" of multiple tests, which are often useful. For example, abnormalities of hepatic function may provide the clue to such nonspecific symptoms as generalized weakness and increased fatigability, suggesting the diagnosis of chronic liver disease. Sometimes a single abnormality, such as an elevated serum calcium level, points to particular diseases, such as hyperparathyroidism or underlying malignancy.

The thoughtful use of screening tests should not be confused with indiscriminate laboratory testing. The use of screening tests is based on the fact that a group of laboratory determinations can be carried out conveniently on a single specimen of blood at relatively low cost. Screening tests are most useful when they are directed towards common diseases or disorders in which the result directs other useful tests or interventions that would otherwise be costly to perform. Biochemical measurements, together with simple laboratory examinations such as blood count, urinalysis, and sedimentation rate, often provide the major clue to the presence of a pathologic process. At the same time, the physician must learn to evaluate occasional abnormalities among the screening tests that may not necessarily connote significant disease. An in-depth workup following a report of an isolated laboratory abnormality in a person who is otherwise well is almost invariably wasteful and unproductive. Among the more than 40 tests that are routinely performed on patients, one or two are often slightly abnormal. If there is no suspicion of an underlying illness, these tests are ordinarily repeated to ensure that the abnormality does not represent a laboratory error. If an abnormality is confirmed, it is important to consider its potential significance in the context of the patient's condition and other test results.

Imaging Techniques The availability of ultrasonography, a variety of scans that employ isotopes to visualize organs heretofore inaccessible, computed tomography, and magnetic resonance imaging has opened new diagnostic vistas and has benefited patients because these new techniques have largely supplanted more invasive ones. While the enthusiasm for noninvasive technology is understandable, the expense entailed in performing these tests is often substantial and should be considered when assessing the potential benefits of the information provided.

PRINCIPLES OF PATIENT CARE Medical Decision-Making Both during and in particular after the physician has taken the history, performed the physical examination, and reviewed the laboratory and imaging data, the challenging process of the differential diagnosis and medical decision-making begins. Formulating a differ-

sincere concern, the willingness to take the time to explain all aspects of the illness, and a nonjudgmental attitude when dealing with patients whose cultures, lifestyles, attitudes, and values differ from those of the physician are just a few of the characteristics of the humane physician. Every physician will, at times, be challenged by patients who evoke strongly negative (or strongly positive) emotional responses. Physicians should be alert to their own reactions to such patients and situations and should consciously monitor and control their behavior so that the patients' best interests remain the principal motivation for their actions at all times.

An important aspect of patient care involves an appreciation of the "quality of life," a subjective assessment of what each patient values most. Such an assessment requires detailed, sometimes intimate knowledge of the patient, which can usually be obtained only through deliberate, unburdened, and often repeated conversations. It is in these situations that the time constraints of a managed care setting may prove problematic.

The famous statement of Dr. Francis Peabody is even more relevant today than when delivered more than three-quarters of a century ago: *"The significance of the intimate personal relationship between physician and patient cannot be too strongly emphasized, for in an extraordinarily large number of cases both the diagnosis and treatment are directly dependent on it. One of the essential qualities of the clinician is interest in humanity, for the secret of the cure of the patient is in caring for the patient."*

CLINICAL SKILLS: History Taking. The written history of an illness should embody all the facts of medical significance in the life of the patient. Recent events should be given the most attention. The patient should, at some point, have the opportunity to tell his or her own story of the illness without frequent interruption and, when appropriate, receive expressions of interest, encouragement, and empathy from the physician. The physician must be alert to the possibility that any event related by the patient, however trivial or apparently remote, may be the key to the solution of the medical problem.

An informative history is more than an orderly listing of symptoms; something is always gained by listening to patients and noting the way in which they describe their symptoms. Inflections of voice, facial expression, gestures, and attitude may reveal important clues to the meaning of the symptoms to the patient. Taking history often involves much data gathering. Patients vary in their medical sophistication and ability to recall facts. Medical history should therefore be corroborated whenever possible. The family and social history can also provide important insights into the types of diseases that should be considered. In listening to the history, the physician discovers not only something about the disease but also something about the patient. The process of history taking provides an opportunity to observe the patient's behavior and to watch for features to be pursued more thoroughly during the physical examination.

The very act of eliciting the history provides the physician with the opportunity to establish or enhance the unique bond that is the basis for the ideal patient-physician relationship. It is helpful to develop an appreciation of the patient's perception of the illness, the patient's expectations of the physician and the medical care system, and the financial and social implications of the illness to the patient. The confidentiality of the patient-physician relationship should be emphasized, and the patient should be given the opportunity to identify any aspects of the history that should not be disclosed.

Physical Examination. Physical signs are objective indications of diseases whose significance is enhanced when they confirm a functional or structural change already suggested by the patient's history. At times, however, the physical signs may be the only evidence of disease.

The physical examination should be performed methodically and thoroughly, with consideration for the patient's comfort and modesty. Although attention is often directed by the history to the diseased organ

or part of the body, the examination of a new patient must extend from head to toe in an objective search for abnormalities. Unless the physical examination is systematic, important segments may be omitted. The results of the examination, like the details of the history, should be recorded at the time they are elicited, not hours later when they are subject to the distortions of memory. Skill in physical diagnosis requires success with experience, but it is not merely technique that determines success in eliciting signs. The detection of a few scattered petechiae, a faint diastolic murmur, or a small mass in the abdomen is not a question of keener eyes and ears or more sensitive fingers but of a mind alert to these findings. Since physical findings are subject to changes, the physical examination should be repeated as frequently as the clinical situation warrants.

Laboratory Tests. The availability of a wide array of laboratory tests has increased our reliance on these studies for the solution of clinical problems. The accumulation of laboratory data does not relieve the physician from the responsibility of careful observation, examination, and study of the patient. It is also essential to bear in mind the limitations of such tests. By virtue of their impersonal quality, confounding factors, and apparent precision, they often gain an aura of authority regardless of the fallibility of the tests themselves, the instruments used in the tests, and the individuals performing or interpreting them. Physicians must weigh the expense involved in the laboratory procedures they order relative to the value of the information they are likely to provide.

Single laboratory tests are rarely ordered. Rather, they are generally obtained as "batteries" of multiple tests, which are often useful for such nonspecific symptoms as generalized weakness and decreased fatigability, suggesting the diagnosis of chronic liver disease. Some points to particular diseases, such as an elevated serum calcium level, may point to particular diseases, such as hyperparathyroidism or underlying malignancy.

The thoughtful use of screening tests should not be confused with indiscriminate laboratory testing. The use of screening tests is based out on the fact that a group of laboratory determinations can be carried out conveniently on a single specimen of blood at relatively low cost. Screening tests are most useful when they are directed towards common diseases or disorders in which the result directs other useful tests or interventions that would otherwise be costly to perform. Biochemical measurements, together with simple laboratory examinations such as blood count, urinalysis, and sedimentation rate, often provide the major clue to the presence of a pathologic process. At the same time, the physician must learn to evaluate occasional abnormalities among the screening tests that may not necessarily connote significant disease. An in-depth workup following a report of an isolated laboratory abnormality in a person who is otherwise well is almost invariably wasteful and unproductive. Among the more than 40 tests that are routinely performed on patients, one or two are often slightly abnormal. If there is no suspicion of an underlying illness, these tests are ordinarily repeated to ensure that the abnormality does not represent a laboratory error. If an abnormality is confirmed, it is important to consider its potential significance in the context of the patient's condition and other test results.

Imaging Techniques. The availability of ultrasonography, a variety of scans that employ isotope to visualize organs heretofore inaccessible, computed tomography, and magnetic resonance imaging has opened new diagnostic vistas and has benefited patients because these new techniques have largely supplanted more invasive ones. While the enthusiasm for noninvasive technology is understandable, the expense entailed in performing these tests is often substantial and should be considered when assessing the potential benefits of the information provided.

PRINCIPLES OF PATIENT CARE. Medical Decision-Making. Both during and in particular after the physical examination, the history, performed the physical examination, and reviewed the laboratory and imaging data, the challenging process of the differential diagnosis and medical decision-making begins. Formulating a differ-

ential diagnosis requires not only a broad knowledge base but also the ability to assess the relative probabilities of various diseases and to understand the significance of missing diagnoses that may be less likely. Arriving at a diagnosis requires the application of the scientific method. Hypotheses are formed, data are collected, and objective conclusions are reached concerning whether to accept or reject a particular diagnosis. Analysis of the differential diagnosis is an iterative process. As new information or test results are acquired, the group of disease processes being considered can be contracted or expanded appropriately. Medical decision-making occurs throughout the diagnostic and treatment process. It involves the ordering of additional tests, requests for consults, and decisions regarding prognosis and treatment. This process requires an in-depth understanding of the natural history and pathophysiology of disease, explaining why these features are strongly emphasized in this textbook. As described below, medical decision-making should be evidence-based, thereby ensuring that patients derive the full benefit of the scientific knowledge available to physicians.

Evidence-Based Medicine. Sackett has defined evidence-based medicine as "the conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients." Rigorously obtained evidence is contrasted with anecdotal experience, which is often biased. Even the most experienced physicians can be influenced by recent experiences with selected patients, unless they are attuned to the importance of using larger, more objective studies for making decisions. The prospectively designed, double-blind, randomized clinical trial represents the "gold standard" for providing evidence regarding therapeutic decisions; but it is not the only source. Valuable evidence about the natural history of disease and prognosis can come from prospective cohort studies and analytic surveys. Persuasive evidence on the accuracy of diagnostic tests can be derived from cross-sectional studies of patients in whom a specific disorder is suspected. Evidence is strengthened immensely when it has been confirmed by multiple investigations, which can be compared with one another and presented in a meta-analysis or systematic overview.

In failing to apply the best and most current evidence, the physician rapid access to the best available evidence is not sufficient for optimal care. The physician must know whether the evidence is relevant to the patient in question and, when it is, the consequences of applying it in any particular situation. The skills and judgment required to apply sound evidence represent an increasing challenge. Indeed, one might redefine a "good doctor" as one who uses the ever-growing body of rigorously obtained evidence (the science of medicine) in a sensible, compassionate manner (the art of medicine).

While an understanding of biologic and physiologic mechanisms forms the basis of contemporary medicine, when a therapeutic modality is selected, the highest priority must often be placed on improving clinical outcome rather than interrupting what is believed to be the underlying process. For example, for decades patients who had suffered myocardial infarction were treated intuitively with drugs that suppress frequent ventricular extrasystoles, since these were believed to be harbingers of ventricular fibrillation and sudden death. Clinical trials, however, have provided firm evidence that the antiarrhythmic agents actually increase the risk of death in such patients. This finding suggests that the extrasystoles are markers of high risk rather than the cause of fatal events.

Practice Guidelines. Physicians are faced with a large, increasing, and often bewildering body of evidence pointing to potentially useful diagnostic techniques and therapeutic choices. The intelligent and cost-effective practice of medicine consists of making selections most appropriate to a particular patient and clinical situation. Professional organizations and government agencies are developing formal clinical practice guidelines in an effort to aid physicians and other caregivers in this endeavor. When guidelines are current and properly applied, they can provide a useful framework for managing patients with particular diagnoses or symptoms. They can protect patients—

particularly those with inadequate health care benefits—from receiving substandard care. Guidelines can also protect conscientious care-givers from inappropriate charges of malpractice and society from the excessive costs associated with the overuse of medical resources. On the other hand, clinical guidelines tend to oversimplify the complexities of medicine. Different groups with differing perspectives may develop divergent recommendations regarding issues as basic as the need for percutaneous transluminal angioplasty in middle-aged persons. Furthermore, guidelines do not—and cannot be expected to—take into account the uniqueness of each individual and of his or her illness. The challenge for the physician is to integrate into clinical practice the useful recommendations offered by the experts who prepare clinical practice guidelines without accepting them blindly or being inappropriately constrained by them.

Assessing the Outcome of Treatment. Clinicians generally use objective and readily measurable parameters to judge the outcome of a therapeutic intervention. For example, findings on physical or laboratory examination—such as the level of blood pressure, the patency of a coronary artery on an angiogram, or the size of a mass on a radiologic examination—can provide information of critical importance. However, patients usually seek medical attention for subjective reasons; they wish to obtain relief from pain, to preserve or regain function, and to enjoy life. The components of a patient's health status or quality of life can include bodily comfort, capacity for physical activity, personal and professional function, sexual function, cognitive function, and overall perception of health. Each of these important areas can be assessed by means of structured interviews or specially designed questionnaires. Such assessments also provide useful parameters by which the physician can judge the patient's subjective view of his or her disability and the response to treatment; particularly in chronic illness. The practice of medicine requires consideration and integration of both objective and subjective outcomes.

Care of the Elderly. Over the next several decades, the practice of medicine will be greatly influenced by the health care needs of the growing elderly population. In the United States the population over age 65 will almost triple over the next 30 years. It is essential that we understand and appreciate the physiologic processes associated with aging, the different responses of the elderly to common diseases, and disorders that occur commonly with aging, such as depression, dementia, frailty, urinary incontinence, and fractures. The elderly have more adverse reactions to drugs, in large part due to altered pharmacokinetics and pharmacodynamics. Commonly used medications such as digoxin and aminoglycosides have prolonged half-lives in the elderly, and tissues such as the central nervous system are more sensitive to certain drugs, such as the benzodiazepines and narcotics. The large number of drugs used by the elderly increases the risk of unwanted interactions, especially when care is provided by several physicians in an uncoordinated manner.

Diseases in Women versus Men. In the past, many epidemiologic studies and clinical trials focused on men. It is now appreciated that there are significant gender differences in diseases that afflict both men and women. Mortality rates are substantially higher in women than in men under the age of 50 suffering acute myocardial infarction. Their male counterparts (and in African-American men in white males); osteoporosis is more common in women, reflecting the menopausal loss of estrogen; diseases involving the immune system, such as lupus erythematosus, multiple sclerosis, and primary biliary cirrhosis, occur more frequently in women; and the average life expectancy of women is greater than that of men. Recently, considerable attention has been paid to women's health issues, a subject that regrettably did not receive sufficient attention in the past. Ongoing study should enhance our understanding of the mechanisms of gender differences in the course and outcome of certain diseases.

Latrogenic Disorders. In an iatrogenic disorder, the deleterious effects of a therapeutic or diagnostic maneuver cause pathology in-

(e.g., the *Candida albicans* protease) have been implicated in fungal invasion of host tissues.

If pathogens are effectively to invade host tissues (particularly the blood), they must avoid the major host defenses represented by complement and phagocytic cells. Bacteria most often avoid these defenses through their cell surface polysaccharides—either capsular polysaccharides or long O-side-chain antigens characteristic of the smooth LPS of gram-negative bacteria. These molecules can prevent the activation and/or depositing of complement opsonins or limit the access of phagocytic cells with receptors for complement opsonins to these molecules when they are deposited on the bacterial surface below the capsule layer. Another potential mechanism of microbial virulence is the ability of some organisms to present the capsule as an apparent self antigen through molecular mimicry. For example, the polysaccharide capsule of group B *N. meningitidis* is chemically identical to an oligosaccharide found on human brain cells.

Immunological studies of capsular polysaccharides have led to an appreciation of the tremendous chemical diversity that can result from the linking of a few monosaccharides. For example, three hexoses can link up in more than 300 different, potentially serologically distinct ways, while three amino acids have only six possible peptide combinations. Capsular polysaccharides have been used as effective vaccines against meningococcal meningitis as well as against pneumococcal and *H. influenzae* infections and may prove to be of value as vaccines against any organisms that express a nontoxic, immunogenic capsular polysaccharide. In addition, most encapsulated pathogens become virulently avirulent when capsule production is interrupted by genetic manipulation; this observation emphasizes the importance of this structure in pathogenesis.

HOST RESPONSE The inflammatory response of the host is critical for interruption and resolution of the infectious process but is also often responsible for the signs and symptoms of disease. Infection promotes a complex series of host responses involving the complement, fibrin, and coagulation pathways. The production of cytokines such as IL-1, TNF- α , and other factors regulated in part by the NF- κ B transcription factor leads to fever, muscle proteolysis, and other effects, as noted above. An inability to kill or contain the microbe usually results in further damage due to the progression of inflammation and infection. For example, in many chronic infections, degranulation of host inflammatory cells can lead to release of host proteases, elastases, histamines, and other toxic substances that can degrade host tissues. Chronic inflammation in any tissue can lead to the destruction of that tissue and to clinical disease associated with loss of organ function, such as sterility from pelvic inflammatory disease caused by chronic infection with *N. gonorrhoeae*.

The nature of the host response elicited by the pathogen often determines the pathology of a particular infection. Local inflammation produces local tissue damage, while systemic inflammation, such as that seen during sepsis, can result in the signs and symptoms of septic shock. The severity of septic shock is associated with the degree of production of host effectors. Disease due to intracellular parasitism results from the formation of granulomas, wherein the host attempts to wall off the parasite inside a fibrotic lesion surrounded by fused epithelial cells that make up so-called multinucleated giant cells. A number of pathogens, particularly anaerobic bacteria, saprophytic, and streptococci, provoke the formation of an abscess, probably because of the presence of zwitterionic surface polysaccharides such as the capsular polysaccharide of *Bacteroides fragilis*. The outcome of an infection depends on the balance between an effective host response that eliminates a pathogen and an excessive inflammatory response that is associated with an inability to eliminate a pathogen and with the resultant tissue damage that leads to disease.

TRANSMISSION TO NEW HOSTS

As part of the pathogenic process, most microbes are shed from the host, often in a form infectious for susceptible individuals. However, the rate of transmissibility may not necessarily be high, even if the disease is severe in the infected individual, as these traits are not linked. Most pathogens exit via the same route by which they entered. Respiratory pathogens by aerosols from sneezing or coughing, through salivary spread, gastrointestinal pathogens by fecal-oral spread, sexually transmitted diseases by venereal spread, and vector-borne organisms by either direct contact with the vector through blood meal or indirect contact with organisms shed into environmental sources such as water. Microbial factors that specifically promote transmission are not well characterized. Respiratory shedding is facilitated by overproduction of mucous secretions, with consequently enhanced sneezing and coughing. Diarrheal toxins such as cholera toxin, *E. coli* heat-labile toxins, and *Shigella* toxins probably facilitate fecal-oral spread of microbial cells in the high volumes of diarrhea fluid produced during infection. The ability to produce phenotypic variants that resist hostile environmental factors (e.g., the highly resistant cysts of *E. histolytica* shed in feces) represents another mechanism of pathogenesis relevant to transmission. Blood parasites such as *Plasmodium* spp. change phenotype after ingestion by a mosquito—a prerequisite for the continued transmission of this pathogen. Venerally transmitted pathogens may undergo phenotypic variation due to the production of specific factors to facilitate transmission, but shedding of these pathogens into the environment does not result in the formation of infective foci.

In summary, the molecular mechanisms used by pathogens to colonize, invade, infect, and disrupt the host are numerous and diverse. Each phase of the infectious process involves a variety of microbial and host factors interacting in a manner that can result in disease. Recognition of the coordinated genetic regulation of virulence factors by the mammalian host emphasizes the complex nature of the host-pathogen interaction. Fortunately, the need for diverse factors in successful infection and disease implies that a variety of therapeutic strategies may be developed to interrupt this process and thereby prevent and treat microbial infections.

BIBLIOGRAPHY

- ANDERSON DM, SCHWENBERG O. Type III molecules of gram-negative pathogens in host defense. *Microbiol Rev* 56:1-10, 1992.
- BECKER EA et al. Chemically responsive HIV-1 coreceptor: Roles in viral entry, pathogenesis, and disease. *Annu Rev Immunol* 17:537, 1999.
- BRANDENBURG TT. T-regulated gene disruption reveals an adhesion indispensable for pathogenicity of *Blastomycosis dermatitidis*. *J Exp Med* 187:1207, 1999.
- CHOW JC et al. Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction. *J Biol Chem* 274:10688, 1999.
- DIVITA DJ et al. The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science* 282:295, 1998.
- EVERT RK et al. How intracellular bacteria survive: Surface modifications that protect resistance to host innate immune responses. *J Infect Dis* 178(Suppl 2):S326, 1999.
- GALL CA et al. Linkage of adhesion, filamentous growth, and virulence in *Candida albicans* to a single gene. *EMBO J* 17:1335, 1998.
- QUINN W, KATZ H-D. Understanding influenza virus pathogenicity. *Trends Microbiol* 7:279-289, 1999.
- MAHONY M. Induction and evasion of host defenses by type 1-piliated uropathogenic *Escherichia coli*. *Science* 284:144, 1999.
- PARA GB et al. Schwann cell type CTR to cause transient epithelial cells. *Nature* 392:79, 1998.
- SCHWENBERG O, ANDERSON DM. Pathogenic aspects of microbes virus infections. *Arch Virol Suppl* 15:139, 1999.
- SOTO OE, HURTADO SI. Bacterial adhesins: Common themes and variations in entry, invasion and assembly. *J Bacteriol* 181:1059, 1999.
- SVANENBERG C et al. Cytokine responses during mucosal infections: Role in disease pathogenesis and host defense. *Curr Opin Microbiol* 2:99, 1999.

Andrew B. Onderdonk

LABORATORY DIAGNOSIS OF INFECTIOUS DISEASES

The laboratory diagnosis of infection requires the demonstration, either direct or indirect, of viral, bacterial, fungal, or parasitic agents in tissues, fluids, or excreta of the host. Clinical microbiology laboratories are responsible for processing these specimens and also for determining the antibiotic susceptibility of bacterial pathogens. Traditional detection of pathogenic agents has relied largely on either the serologic visualization of pathogens in clinical material or the growth of microorganisms in the laboratory. Identification is generally based on phenotypic characteristics, such as fermentation profiles for bacteria, cytopathic effects in tissue culture for viral agents, and microscopic morphology for fungi and parasites. These techniques are reliable but are often time-consuming. Increasingly, the use of nucleic acid probes is becoming a standard detection and/or identification method in the clinical microbiology laboratory, gradually replacing serotypic characterization and microscopic visualization methods.

DETECTION METHODS

Examination of the methods employed in the clinical microbiology laboratory has led to the development of strategies for detection of pathogenic agents through nonvisual biologic signal detection systems. Much of this methodology is based on either computerization of detection systems with relatively inexpensive but sophisticated computation or the use of nucleic acid probes directed at specific DNA or RNA targets. This chapter discusses both the methods that are currently available and those that are being developed.

BIOLOGIC SIGNALS A biologic signal is a material that can be serologically differentiated from other substances present in the same physical environment. Key issues in the use of a biologic (or chemical) signal are distinguishing it from background noise and translating it into meaningful information. Examples of biologic signals applicable to clinical microbiology include structural components of bacteria, fungi, and viruses; specific antigens; membrane and products; unique DNA or RNA base sequences; enzymes; toxins; or other proteins and surface polysaccharides.

DETECTION SYSTEMS A detector is used to sense a signal and to discriminate between the signal and background noise. Detection systems range from the trained eyes of a microbiologist assessing morphologic variations to sensitive electronic instruments, such as gel electrophoresis coupled to computer systems for signal analysis. The sensitivity with which signals can be detected varies widely. It is essential to use a detection system that discerns small amounts of signal even when biologic background noise is present—i.e., that is both sensitive and specific. Common detection systems include immunofluorescence, chemiluminescence for DNA/RNA probes, flame ionization detection of short- or long-chain fatty acids, and detection of substrate utilization or end-product formation as color changes, of enzyme activity as a change in light absorbance, of turbidity changes, of cytopathic effects in cell lines, and of particle agglutination.

AMPLIFICATION Amplification enhances the sensitivity with which weak signals can be detected. The most common microbiologic amplification technique is growth of a single bacterium into a discrete colony on an agar plate or into a suspension containing many identical organisms. The advantage of growth as an amplification method is that it requires only an appropriate growth medium; the disadvantage is that it requires only an appropriate growth medium. More rapid specific amplification of biologic signals can be achieved with techniques such as polymerase chain reactions (PCRs, for DNA/RNA), enzyme immunoassays (EIAs, for antigens and antibodies), electronic amplification (for gas-liquid chromatography assays), antibody capture methods (for concentration and/or separation), and selective filtration

or centrifugation. Although a variety of methods are available for the amplification and detection of biologic signals in research, thorough testing is required before they are validated as diagnostic assays.

DIRECT DETECTION

MICROSCOPY The field of microbiology has been defined largely by the development and use of the microscope. The examination of specimens by microscopic methods readily provides useful diagnostic information. Staining techniques permit organisms to be seen more clearly.

The simplest method for microscopic evaluation is the wet mount, which is used, for example, to examine cerebrospinal fluid (CSF) for the presence of *Cryptococcus neoformans*, with India ink as a background against which to visualize large-encapsulated yeast cells. Wet mounts with dark-field illumination are also used to detect spirochetes from genital lesions and to reveal *Borrelia* or *Leptospira* in blood. Skin scrapings and hair samples can be examined with use of either 10% KOH wet-mount preparations or the calcofluor white method and ultraviolet illumination to detect fungal elements as fluorescing structures. Staining of wet mounts—for example, with lactophenol cotton blue stain for fungal elements—is often used for morphologic identification. These techniques enhance signal detection and decrease the background, making it easier to identify specific fungal structures.

STAINING Gram's Stain: Without staining, bacteria are difficult to see at the magnifications (400 to 1000 \times) used for their detection. Although simple one-step stains can be used, differential stains are more common. Gram's stain differentiates between organisms with thick peptidoglycan cell walls (gram-positive) and those with outer membranes that can be dissolved with alcohol or acetone (gram-negative).

Gram's stain is particularly useful for examining sputum for polymorphonuclear leukocytes (PMNs) and bacteria. Sputum specimens with 25 or more PMNs and fewer than 10 epithelial cells per low-power field often provide clinically useful information. However, the presence in "sputum" samples of more than 10 epithelial cells per low-power field and of multiple bacterial types suggests contamination with oral microflora. Despite the difficulty of discriminating between normal microflora and pathogens, Gram's stain may prove useful for specimens from areas with a large resident microflora if a useful biologic marker (signal) is available. Gram's staining of vaginal swab specimens can be used to detect epithelial cells covered with gram-positive bacteria in the absence of lactobacilli and the presence of gram-negative rods—a scenario regarded as a sign of bacterial vaginosis. Similarly, examination of stained stool specimens for leukocytes is useful as a screening procedure before testing for *Clostridium difficile* toxin or other enteric pathogens.

The examination of CSF and joint, pleural, or peritoneal fluid with Gram's stain is useful for determining whether bacteria and/or PMNs are present. The sensitivity is such that $>10^6$ bacteria per milliliter should be detected. Centrifugation is often performed before staining to concentrate specimens thought to contain low numbers of organisms. The pellet is examined after staining. This simple method is particularly useful for examination of CSF for bacteria and white blood cells or of sputum for acid-fast bacilli (AFB).

Acid-Fast Stain The acid-fast stain identifies organisms that retain carbol fuchsin dye after acid/organic solvent disruption (e.g., *Mycobacterium* spp.). Modifications of this procedure allow the differentiation of *Acetomyces* from *Nocardia* or other weakly acid-fast organisms. The acid-fast stain is applied to sputum, other fluids, and tissue samples when AFB (e.g., *Mycobacterium* spp.) are suspected. The identification of the pink/red AFB against the blue background of the counterstain requires a trained eye, since few AFB may be detected in an entire smear, even when the specimen has been concentrated by centrifugation. An alternative method is the auramine-rhodamine combination fluorescent dye technique.

SPECIMEN COLLECTION AND TRANSPORT To culture bacterial, mycotic, or viral pathogens, an appropriate sample must be placed into the proper medium for growth (amplification). The success of efforts to identify a specific pathogen often depends on the collection and transport process coupled to a laboratory-processing algorithm suitable for the specific sample type. In some instances, it is better for specimens to be placed at the time of collection rather than being first being transported to the laboratory (e.g., urethral swabs being cultured for *Neisseria gonorrhoeae* or sputum specimens for pneumococci). In general, the more rapidly a specimen is placed onto a transport medium, the better the chance for isolating bacterial pathogens. Appendix B lists procedures for collection and transport of common specimens.

The detection of microbial pathogens in blood is difficult because of the small number of organisms present in the sample is often low and because of the high degree of background fluorescence from the blood components. The number of organisms present in the sample is often low and the background fluorescence from the blood components is high. The number of organisms present in the sample is often low and the background fluorescence from the blood components is high. The number of organisms present in the sample is often low and the background fluorescence from the blood components is high.




FIGURE 121-1 Common specimen-processing algorithms used in clinical microbiology laboratories. Abbreviations: BAP, blood agar plate; CMV, cytochrome oxidase; CPE, cytopathic effects; CSF, cerebrospinal fluid; DFA, direct fluorescent antibody; EIA, enzyme immunoassay; ESBL, extended-spectrum β -lactamase; GBS, group B *Streptococcus*; GC, *Neisseria gonorrhoeae*; HACEC, gram-liquid chromatase; HACEC, *Haemophilus aphrodisiac* (GLC, gas-liquid chromatography).

Automated systems also have been applied to the detection of microbial growth from specimens other than blood, such as peritoneal and other normally sterile fluids. *Mycobacterium* spp. can be detected in certain automated systems if appropriate liquid media are used for

Measurement of serum antibody provides an indirect marker for past or current infection with a specific viral agent or other pathogens, including *Brucella*, *Legionella*, *Rickettsia*, and *Helicobacter pylori*. The biologic signal is usually either IgM or IgG antibody directed at surface-expressed antigens(s). The detection systems include those used for bacterial antigens (agglutination reactions, immunofluorescence, and ELISA) and unique systems such as hemolysis inhibition and complement fixation. Serologic methods generally fall into two categories: those that determine protective antibody levels and those that measure changing antibody titers during infection. Determination of an antibody response as a measure of current immunity is important in the use of viral agents for which there are vaccines, such as rubella virus or varicella-zoster virus; assays for this purpose normally use one or two dilutions of serum for a qualitative determination of protective

appearance. Broth tubes containing specific substrates are commonly employed for phenotypic characterization. While such methods have been used since the time of Pasteur, their simplicity and low cost continue to make them appealing today.

CLASSIC PHENOTYPING Automated systems allow rapid phenotypic identification of bacterial pathogens. Most such systems are based on biotyping techniques, in which isolates are grown on multiple substrates and the reaction patterns is compared with known patterns for various bacterial species. This procedure is relatively fast, and commercially available systems include miniaturized fermentations, coding to simplify recording of results, and probability calculations for the most likely pathogen. If the biotyping approach is automated and the reading process is coupled to computer-based data analysis, rapidly growing organisms, such as *Enterobacteriaceae*, can be identified within hours of detection on agar plates.

Several systems use preformed enzymes for even speedier identification (within 2 to 3 h). Such systems do not rely on bacterial growth per se to determine whether a substrate has been used or not. They employ a heavy inoculum in which specific bacterial enzymes are present in sufficient quantity to convert substrate to product rapidly. In addition, some systems use fluorogenic substrate-lead-product detection methods to increase sensitivity (through signal amplification).

GAS-LIQUID CHROMATOGRAPHY Gas-liquid chromatography is often used to detect metabolic end products of bacterial fermentations. One common application is identification of short-chain fatty acids produced by obligate anaerobes during glucose fermentation. Because the types and relative concentrations of volatile acids differ among the various genera and species that make up this group of organisms, such information serves as a metabolic "fingerprint" for a particular isolate.

Gas-liquid chromatography can be coupled to a sophisticated signal-analyzer software system for identification and quantitation of long-chain fatty acids (LCFAs) in the outer membrane and cell walls of bacteria and fungi. For many given species, the types and relative concentrations of LCFAs are distinctive enough to allow identification of even closely related species. An organism may be identified definitively within a few hours after detection of growth on appropriate media. LCFA analysis is one of the most advanced procedures currently available for phenotypic characterization.

NUCLEIC ACID PROBES Techniques for the detection and quantitation of specific DNA and RNA base sequences in clinical specimens have become powerful tools for the diagnosis of bacterial, viral, parasitic, and fungal infections. The basic strategy is to detect a relatively short sequence of bases specific for a particular pathogen on single-stranded DNA or RNA by hybridization of a complementary sequence of bases (probe) coupled to a "reporter" system that serves as the signal for detection. Detection of an organism by nucleic acid probes offers a decided advantage over culture methods for difficult-to-grow organisms. Current technology encompasses a wide array of methods for amplification and signal detection, some of which have been approved by the U.S. Food and Drug Administration (FDA) for clinical diagnosis.

Use of nucleic acid probes generally involves lysis of intact cells and denaturation of the DNA or RNA to render it single-stranded. The probe may be hybridized to the target sequence in a solution or on a solid support, depending on the system employed. In situ hybridization of a probe to a target is also possible and allows the use of probes with agents present in tissue specimens. Once the probe has been hybridized to the target (biologic signal), a variety of strategies may be employed to amplify and/or quantify the target-probe complex (Fig. 121-2).

Probes for Direct Detection of Pathogens in Clinical Specimens Nucleic acid probes are available commercially for direct detection of various bacterial and parasitic pathogens, including *L. pneumophila*, *Chlamydia trachomatis*, *M. gonorrhoeae*, group A *Streptococcus*, *Giardia lamblia*, *Mycobacterium hominis*, and *Giardia lamblia*. In addition, probes for direct detection of human papillomavirus, *Con-*

trichomonas vaginalis have been approved. An assortment of probes for confirming the identity of cultured pathogens such as *Mycobacterium* and *Salmonella* spp. are also available. Probes for the direct detection of bacterial pathogens are often aimed at highly conserved 16S ribosomal RNA sequences, of which there are many more copies than there are of any single genomic DNA sequence in a bacterial cell. The sensitivity and specificity of probe assays for direct detection are comparable to those of more traditional assays, including EIA and culture. Many laboratories have developed their own probes for pathogens; however, unless a method-validation protocol for diagnostic testing has been performed, the use of such probes is restricted to research by federal law in the United States.

Nucleic Acid Probe Target-Amplification Strategies In theory, a single target nucleic acid sequence can be amplified to detectable levels. There are several strategies for target and/or probe amplification, including PCR, ligase chain reaction, strand displacement amplification, and self-sustaining sequence replication. In each case, target sequence or hybridized probe is amplified exponentially to obtain sufficient signal for detection, usually by the attachment of chemiluminescent reporter groups to the amplified product. The PCR strategy requires repeated heating of the DNA or RNA to separate the two complementary strands of the double helix, hybridization of a primer sequence to the appropriate target sequence, target amplification using the PCR for complementary strand extension, and signal detection via a labeled probe. The sensitivity of such assays is far greater than that of traditional assay methods such as culture. However, the care with which the assays are performed is important, because cross-contamination of clinical material with DNA or RNA from other sources (even at low levels) can cause false-positive results. An alternative method employs transcription-mediated amplification, in which an RNA target sequence is converted to DNA, which is then exponentially transcribed into RNA target. The advantage of this method is that only a single heating/annealing step is required for amplification. At present, signal amplification assays for *Mycobacterium tuberculosis*, *N. gonorrhoeae*, *C. trachomatis*, and *M. hominis* are on the market. Again, many laboratories have used commercially available *in-house* polymerase, probe sequences, and reagents to develop "in-house" assays for diagnostic use. Issues related to quality control, interpretation of results, sample processing, and regulatory requirements have slowed the commercial development of diagnostic assay kits.

Signal Amplification Strategies Alternative systems for signal amplification have great appeal, particularly for quantitative determination of the amount of target present in a given specimen. With the advent of newer therapeutic regimens for HIV-associated disease, cytomegalovirus infection, and hepatitis C virus infection, the response to therapy has been monitored by determining both genotype and "viral load" at various times after treatment initiation. Target amplification (PCR, transcription-mediated amplification) is difficult to control in a manner that allows accurate determination of the original target (genome) concentration. In other systems, probes attached to complementary target sequences are amplified by the attachment of a second probe and an amplification malimase to the original probe. In one such system, branched-chain DNA (bDNA)-based amplification, bDNA is attached to a site different from the target-binding sequence of the original probe. Chemiluminescent-labeled oligonucleotides can then bind to multiple repeating sequences on the bDNA. The amplified bDNA signal is detected by chemiluminescence. Alternatively, a DNA probe may be attached to an RNA target and the resulting DNA/RNA hybrid captured on a solid support by antibody specific for chemiluminescent hybrids (concentration/amplification) and detected by chemiluminescence. Another antibody specific for DNA/RNA hybrids. Both methods can be used to determine the approximate number of target copies (viral) in the starting material. The advantage of these systems over PCR is that only a single heating/annealing step is required to hybridize the target-binding probe to the target sequence for amplification. Application of Nucleic Acid Probe Technology Nucleic acid probe technology is being used to identify difficult-to-grow or noncultivable bacterial pathogens, such as *Mycobacterium*, *Legionella*




FIGURE 121-2 Strategies for amplification and/or detection of a target probe. DNA or RNA extracted from microorganisms is heated to create single-stranded (ss) DNA/RNA containing appropriate target sequences. These target sequences may be hybridized directly (direct detection) with probes attached to reporter molecules; they may be amplified by repetitive cycles of complementary strand extension (polymerase chain reaction) before attachment

of a reporter probe, or the original target-probe signal may be amplified via hybridization with an additional probe containing multiple copies of a second reporter target sequence (branched-chain DNA or bDNA). bDNA/RNA hybrids can also be "captured" on a solid support (hybrid capture), with antibody directed at the DNA/RNA hybrids used to concentrate them and a second antibody coupled to a reporter molecule attached to the captured hybrid.

screen for infection control problems, such as methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant *Enterococcus faecium*, or extended-spectrum β -lactamase-producing organisms. Two approaches are useful. The first is a qualitative assessment of susceptibility, with responses categorized as susceptible, resistant, or intermediate. This approach can involve either the placement of paper disks containing antibiotics on an agar surface inoculated with the bacterial strain to be tested (Kirby-Bauer or disk/diffusion method), with measurement of the zones of growth inhibition following incubation, or the use of broth tubes containing a set concentration of antibiotic (breakpoint method). These methods have been carefully calibrated against quantitative methods and clinical experience with each antibiotic, and zones of inhibition and breakpoints have been calculated on a species-by-species basis.

The second approach is to inoculate the test strain of bacteria into a series of broth tubes (or agar plates) with increasing concentrations of antibiotic. The lowest concentration of antibiotic that inhibits microbial growth in this test system is known as the *minimum inhibitory concentration* (MIC). If tubes in which no growth occurs are subcultured, the minimum concentration of antibiotic required to kill the starting inoculum can also be determined (*minimum bactericidal concentration*, or MBC). Quantitative susceptibility testing by the microbroth dilution technique, a miniaturized version of the broth dilution technique using microtiter plates, lends itself to automation and is commonly used in larger clinical laboratories.

Brucella, *Rickettsia*, *Babesia*, *Borrelia*, and *Tropheryma whippelii*. Amplification methods are also being used to detect chronic viral infections, such as herpes simplex encephalitis, cytomegalovirus infection, and hepatitis C. The monitoring of therapy with quantitative viral load testing is a significant new application of nucleic acid technology. Further applications will likely include the replacement of culture for identification of many pathogens with solid-state DNA/RNA chip technology, in which thousands of unique nucleic acid sequences can be detected on a single computer chip. Probe technology also has the potential to detect viral pathogens faster than is possible with current culture techniques. However, if laboratories are to take full advantage of probe technology, the cost of reagents and assay automation must be competitive with the cost of existing methodology. At present, the detection of agents such as *C. trachomatis* or *N. gonorrhoeae* by probe technology is more expensive for most laboratories than detection by traditional culture or EIA. Moreover, because automated processing equipment is just beginning to find its way into the laboratory for these assays, nucleic acid amplification methods are both more labor-intensive and more expensive than other detection systems. In the absence of clear documentation of clinical utility, many laboratories continue to wait for FDA approval of commercially available DNA/RNA probe assays rather than validating in-house assays.

CLINICAL MICROBIOLOGY TESTING A principal responsibility of the clinical microbiology laboratory is to determine which antimicrobial agents inhibit a specific bacterial isolate. Such testing is used to

EDITORS OF PREVIOUS EDITIONS

T. R. HARRISON Editor-in-Chief, Editions 1, 2, 3, 4, 5	R. G. PETERSDORF Editor, Editions 6, 7, 8, 9, 11, 12, 13 Editor-in-Chief, Edition 10
W. R. RESNICK Editor, Editions 1, 2, 3, 4, 5	J. D. WILSON Editor, Editions 9, 10, 11, 13, 14 Editor-in-Chief, Edition 12
M. M. WINTROBE Editor, Editions 1, 2, 3, 4, 5 Editor-in-Chief, Editions 6, 7	J. B. MARTIN Editor, Editions 10, 11, 12, 13, 14
G. W. THORN Editor, Editions 1, 2, 3, 4, 5, 6, 7 Editor-in-Chief, Edition 8	A. S. FAUCI Editor, Editions 11, 12, 13 Editor-in-Chief, Edition 14
R. D. ADAMS Editor, Editions 2, 3, 4, 5, 6, 7, 8, 9, 10	R. ROOT Editor, Edition 12
P. B. BEESON Editor, Editions 1, 2	D. L. KASPER Editor, Editions 13, 14
I. L. BENNETT, JR. Editor, Editions 3, 4, 5, 6	S. L. HAUSER Editor, Edition 14
E. BRAUNWALD Editor, Editions 6, 7, 8, 9, 10, 12, 13, 14 Editor-in-Chief, Edition 11	D. L. LONGO Editor, Edition 14
K. J. ISSELBACHER Editor, Editions 6, 7, 8, 10, 11, 12, 14 Editor-in-Chief, Editions 9, 13	

HARRISON'S 15TH EDITION

PRINCIPLES OF INTERNAL MEDICINE

EDITORS

EUGENE BRAUNWALD, MD, MD(Hon), ScD(Hon) Distinguished Hersey Professor of Medicine, Faculty Dean for Academic Programs at Brigham and Women's Hospital and Massachusetts General Hospital, Harvard Medical School; Vice-President for Academic Programs, Partners HealthCare Systems, Boston	STEPHEN L. HAUSER, MD Betsy Auker Life Professor and Chairman, Department of Neurology, University of California San Francisco, San Francisco
ANTHONY S. FAUCI, MD, ScD(Hon) Chief, Laboratory of Immunoregulation; Director, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda	DAN L. LONGO, MD Scientific Director, National Institute on Aging, National Institutes of Health, Bethesda and Baltimore
DENNIS L. KASPER, MD, MA(Hon) William Ellery Channing Professor of Medicine, Professor of Microbiology and Molecular Genetics, Executive Dean for Academic Programs, Harvard Medical School; Director, Channing Laboratory, Department of Medicine, Brigham and Women's Hospital, Boston	J. LARRY JAMESON, MD, PhD Irving S. Cutter Professor and Chairman, Department of Medicine, Northwestern University Medical School; Physician-in-Chief, Northwestern Memorial Hospital, Chicago

McGraw-Hill

MEDICAL PUBLISHING DIVISION

New York	San Francisco	Washington, DC	Auckland	Bogotá	Caracas	Lisbon	London
Madrid	Mexico City	Milan	Montreal	New Delhi	San Juan	Singapore	Sydney
Tokyo	Toronto						



Note: Dr. Fauci and Dr. Longo's works as editors and authors were performed outside the scope of their employment as U.S. government employees. These works represent their personal and professional views and not necessarily those of the U.S. government.

**Harrison's
PRINCIPLES OF INTERNAL MEDICINE
Fifteenth Edition**

Copyright © 2001, 1998, 1994, 1991, 1987, 1983, 1980, 1977, 1974, 1970, 1966, 1962, 1958 by The McGraw-Hill Companies, Inc. All rights reserved. Printed in the United States of America. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a data base or retrieval system, without the prior written permission of the publisher.

1234567890 DOWDOW 0987654321

ISBN 0-07-007272-8 (Combo)

0-07-007273-6 (Vol. 1)

0-07-007274-4 (Vol. 2)

0-07-913686-9 (Set)

FOREIGN LANGUAGE EDITIONS

Arabic (Thirteenth Edition)—McGraw-Hill Libri Italia srl (est. 1996)

Chinese (Twelfth Edition)—McGraw-Hill Book Company—Singapore © 1994

Croatian (Thirteenth Edition)—Placebo, Split, Croatia

French (Fourteenth Edition)—McGraw-Hill Publishing Co., Maidenhead, UK © 1999

German (Fourteenth Edition)—McGraw-Hill Publishing Co., Maidenhead, UK © 1999

Greek (Fourteenth Edition)—Parissianos, Athens, Greece © 2000

Italian (Fourteenth Edition)—McGraw-Hill Libri Italia srl, Milan © 1999

Japanese (Eleventh Edition)—Hirokawa © 1991

Polish (Fourteenth Edition)—Czelej Publishing Company, Lubin, Poland (est. 2000)

Portuguese (Fourteenth Edition)—McGraw-Hill Interamericana do Brasil Ltda © 1998

Romania (Fourteenth Edition)—Teora Publishers, Bucharest, Romania (est. 2000)

Spanish (Fourteenth Edition)—McGraw-Hill Interamericana de Espana, Madrid © 1998

Turkish (Thirteenth Edition)—McGraw-Hill Libri Italia srl (est. 1996)

This book was set in Times Roman by Progressive Information Technologies. The editors were Martin Wonsiewicz and Mariapaz Ramos Englis. The production director was Robert Laffler. The index was prepared by Irving C. Tullar. The text and cover designer was Marsha Cohen/Parallelogram Graphics.

R. R. Donnelley and Sons, Inc. was the printer and binder.

Library of Congress Cataloging-in-Publication Data

Harrison's principles of internal medicine—15th ed./editors, Eugene Braunwald . . . [et al.] p. cm.

Includes bibliographical references and index.

ISBN 0-07-913686-9 (set)—ISBN 0-07-007273-6 (v. 1)—ISBN 0-07-0072744-4 (v. 2)

1. Internal medicine. I. Braunwald, Eugene, date

RC46.H333 2001

616—dc21

00-063809



[Short Contents](#) | [Full Contents](#)

[Other books @ NCBI](#)

Navigation

About this book

**Appendix I.
Immunologists'
Toolbox**

Immunization.

The detection, measurement, and characterization of antibodies and their use as research and diagnostic tools.

➔ Isolation of lymphocytes.

Characterization of lymphocyte specificity, frequency, and function.

Detection of immunity *in vivo*.

Manipulation of the immune system.

Immunobiology ➔ Appendix I. Immunologists' Toolbox ➔
Isolation of lymphocytes.




Figure A.23. Peripheral blood mononuclear cells can be isolated from whole blood by Ficoll-Hypaque™ centrifugation. Diluted anticoagulated blood (left panel) is layered over Ficoll-Hypaque™ and centrifuged. Red blood cells and polymorphonuclear leukocytes or granulocytes are more dense and centrifuge through the Ficoll-Hypaque™, while mononuclear cells consisting of lymphocytes together with some monocytes band over it and can be recovered at the interface (right panel).

Search

☒ This book ☐ All books

☐ PubMed

© 2001 by Garland Science

Consequences of different diagnostic 'gold standards' in test accuracy research: Carpal Tunnel Syndrome as an example

Lucas M Bachmann,^{1,2} Peter Jüni,^{1,3,4*} Stephan Reichenbach,^{1,3,4} Hans-Rudolf Ziswiler,³
 Alfons G Kessels^{2,5} and Esther Vögelin⁶

Accepted 20 April 2005

Test accuracy studies assume the existence of a well-defined illness definition and clear-cut diagnostic gold standards or reference standards. However, in clinical reality illness definitions may be vague or a mere description of a set of manifestations, mostly clinical signs and symptoms. This can lead to disagreements among experts about the correct classification of an illness and the adequate reference standard. Using data from a diagnostic accuracy study in carpal tunnel syndrome, we explored the impact of different definitions on the estimated test accuracy and found that estimated test performance characteristics varied considerably depending on the chosen reference standard. In situations without a clear-cut illness definition, randomized controlled trials may be preferable to test accuracy studies for the evaluation of a novel test. These studies do not determine the diagnostic accuracy, but the clinical impact of a novel test on patient management and outcome.

Keywords Sensitivity and specificity, ROC curve, reference standards, carpal tunnel syndrome, ultrasonography

The notion of a diagnostic gold standard or reference standard pertains to the best available method for establishing the presence or absence of a condition of interest,¹ i.e. the independent and correct classification of what is meant to be the illness.² The traditional concept of a reference standard depends on a high level of biological understanding of the target condition and its causal underlying mechanisms. Typically, a morphological verification such as histopathology or angiography, is used to establish a 'definite diagnosis'. This definite diagnosis is assumed to be a reasonably reliable proxy measure of the true presence or absence of the condition of interest.

In conventional diagnostic accuracy studies, the usefulness of a novel test for the inclusion or exclusion of a specific condition will be determined by comparing the results of the test with the definite diagnosis ascertained by the reference standard.

However, in clinical reality the biological understanding of conditions is frequently unclear. Illness definitions are vague or a mere description of a set of manifestations. In fields such as psychiatry and rheumatology, clinicians frequently use 'syndromal diagnoses' consisting of a characteristic pattern of signs and symptoms,³ while the biological understanding of the condition, of its causes, and its manifestations is incomplete and there is controversy about the manifestations that have to be combined to ensure accurate representation of the condition. In other situations, the biological understanding of the condition may be comprehensive, but the measurement of signs or symptoms is inaccurate.

Two extreme conceptualizations of the reference standard may implicitly or explicitly be used in such circumstances. One extreme ignores potential controversies and assumes a well-defined illness, which is objectively and reproducibly represented by the outcome of one or several laboratory tests. The other extreme ignores potentially useful biological measures and focuses exclusively on patient outcomes or on the need for an intervention. While these two outlooks aim at describing the same issue, they may create a schism when evaluating a diagnostic test. Below, we will explore this in a clinical example of an accuracy study previously published by our group in the field of rheumatology⁴ and discuss the potential implications for clinical research into conditions without a clear-cut reference standard by which to establish a diagnosis.

¹ Department of Social and Preventive Medicine, University of Berne, Switzerland.

² Herten Centre, University of Zurich, Switzerland.

³ Department of Rheumatology and Clinical Immunology, Inselspital University of Berne, Switzerland.

⁴ MRC Health Services Research Collaboration, Department of Social Medicine, University of Bristol, UK.

⁵ Department of Clinical Epidemiology and Medical Technology Assessment, Maastricht University Hospital, Maastricht, The Netherlands.

⁶ Department of Hand Surgery, Inselspital, University of Berne, Switzerland.

* Corresponding author. E-mail: juni@ispim.unibe.ch

Clinical example

Carpal Tunnel Syndrome (CTS) is an important cause of functional impairment and pain of the hand, which presumably results from a compression of the median nerve at the wrist. Unfortunately, there is no universally accepted reference standard to establish the diagnosis. In our experience, two different approaches towards CTS classification are used. Neurologists traditionally establish the definite diagnosis based more on the outcome of nerve conduction studies than on the patients' signs and symptoms. In contrast, hand surgeons appear to give considerably more importance to the patients' signs and symptoms, the severity of complaints and the likely need for and success of a surgical intervention than to nerve conduction studies when establishing the definite diagnosis. In our accuracy study,⁴ we relied on current practice and pre-specified the neurologists' definite diagnosis as the reference standard. Here, we determine the impact of using either of the two 'reference standards' on the estimated test accuracy of sonography in patients with suspected CTS.

Methods and results

Details of methods are reported elsewhere.⁴ We assessed 77 patients for eligibility, excluded 3 because of traumatic wrist lesions, and enrolled 74 referred to the outpatient clinic of the Department of Hand Surgery at the University Hospital Berne, Switzerland, between January and December 2002.

Patients included in the study had a mean age of 51 years and 48 were females (65%). The flow of patients through the various stages of the study is described elsewhere.⁴ Essentially, 101 wrists from 71 patients were included in the analysis.

Standardized nerve conduction studies were performed by one of several neurologists, who were unaware of the results of the sonographic examination. The sonographic evaluations were performed by a rheumatologist experienced in musculoskeletal sonography, who was unaware of the results of the nerve conduction studies and of the patients' signs and symptoms. He performed transverse imaging of the median nerve for the area ranging from the distal forearm to the outlet of the carpal tunnel and measured the largest cross-sectional area of the median nerve in square millimetres. We used this measure as a single diagnostic indicator, assuming that an increase in cross-sectional areas is associated with an increasing likelihood of disease or disease severity.

Table 1 presents a comparison of definite diagnoses according to neurologists' and hand surgeons' judgements. Overall agreement was 86%. One out of 23 wrists classified as normal by the neurologists was considered as CTS by the hand surgeons (4%). This wrist had normal nerve conduction studies.

Table 1 2 × 2 contingency table comparing reference standard classifications according to neurologists and hand surgeons

	Hand surgeons' judgements		Total
	CTS present	CTS absent	
Neurologists' judgements			
CTS present	65	13	78
CTS absent	1	22	23
Total	66	35	101

Conversely, 13 out of 78 wrists classified as CTS by the neurologists were considered normal by the hand surgeons (17%); all 13 wrists had pathological nerve conduction studies. The resulting kappa for the agreement between the two illness definitions was 0.67 [95% confidence interval (CI) 0.48–0.85].

For both reference standards, we fitted a receiver operating characteristic (ROC) curve for diagnosis of CTS by sonography, using a maximum likelihood logistic regression model based on robust standard errors, which allowed for the correlation of characteristics of wrists within patients and compared the area under the ROC curve. Figure 1 shows the fitted ROC curves using either the neurologists' judgements (top) or the hand surgeons' judgements (bottom) as the reference standard. The area under the ROC curve for ultrasound was 0.89 based on neurologists' judgements (95% CI 0.82–0.96) and 0.77 based on hand surgeons' judgements (95% CI 0.68–0.87). The difference between the two areas under the ROC curve was 0.12 (95% CI 0.0–0.23).




Figure 1 Fitted ROC curves (solid curve) for diagnosis of CTS by sonography with 95% confidence interval (dotted curves), considering the neurologists' definite diagnosis (top) or the hand surgeons' definite diagnosis as the reference standard (bottom). The broken diagonal line represents a hypothetical ROC curve of a test that yields no diagnostic information

Discussion

Even though the agreement between the two employed illness definitions was substantial (a kappa of 0.67), the estimated test performance of ultrasound varied considerably depending on the definition used as the reference standard. The diagnostic accuracy of sonography in patients with suspected CTS was good to excellent according to one reference standard but only moderate according to the other.

The lack of consensus on an illness definition may impede a valid evaluation of diagnostic technology in test accuracy studies. Considering that the final purpose of any novel test is to improve patient management and outcome, the traditional paradigm of test accuracy studies will only be useful if a reference standard is chosen that either has a strong association with patient outcome or a direct relationship with patient management. In our accuracy study⁴ we argued, for example, that the neurologists' definite diagnosis directly pertains to clinical decision making and patient management.

Ultimately, the use of a diagnostic test and its potential therapeutic consequences can be considered as two consecutive steps of the same management strategy. Analogous to traditional research into therapeutic interventions, randomized trials may be designed to compare different strategies. In such trials, patients will be randomly allocated to a management

strategy that includes the use of a novel test under evaluation, or to a strategy that uses standard tests only. Ascertained outcomes may relate to parameters of patient management (e.g. length of hospital stay), to patient outcome (e.g. pain), or to the total cost of management per patient.⁵ If an unanimously accepted reference standard is lacking, as is the case in CTS, such randomized controlled trials may be more appropriate than test accuracy studies to determine the usefulness of a novel diagnostic test.

References

- ¹ Bossuyt PM, Reitsma JB, Bruns DE, *et al.* Towards complete and accurate reporting of studies of diagnostic accuracy: The STARD initiative. *Ann Intern Med* 2003;138:40-44.
- ² Wulff HR, Goetzsche PC. Diagnosis. In: *Rational diagnosis and treatment: evidence-based clinical decision making*. Third Edition. Oxford: Blackwell Publishing Ltd, 2000, p. 67.
- ³ Wulff HR, Goetzsche PC. The disease classification. In: *Rational diagnosis and treatment: evidence-based clinical decision making*. Third Edition. Oxford: Blackwell Publishing Ltd, 2000, p. 39.
- ⁴ Ziswiler HR, Reichenbach S, Vögelin F, Bachmann LM, Villiger PM, Jüni P. Diagnostic value of sonography in patients with suspected carpal tunnel syndrome: a prospective study. *Arthritis Rheum* 2005;52:304-11.
- ⁵ Bossuyt PM, Lijmer JG, Mol BW. Randomised comparisons of medical tests: sometimes invalid, not always efficient. *Lancet* 2000;356:1844-47.